

# WEST AFRICAN JOURNAL OF ORTHODONTICS

VOLUME 8, NUMBER 2

ISSN 2315-9502

DECEMBER 2019

**Dental Arch Dimensions in Sickle  
Cell Anaemia**



**Tooth Movement in the resolution  
of Crowding**



**Oral Ornaments and Orthodontic  
Treatment**



**Management of Midline Diastema**



**Orthodontic Management of  
Anterior Crowding**

# Orthodontic Management of Maxillary And Mandibular Midline Diastema

Umeh OD<sup>a</sup>, Offojebe UL<sup>b</sup>

## Abstract

A 23 year-old male with the chief complaint of upper and lower central gaps in his teeth. Orthodontic evaluation gave a diagnosis of Angle's Class I malocclusion on Skeletal Pattern 2 with maxillary and mandibular midline diastema, upper and lower anterior spacing, rotated teeth, abnormal frenal attachment, and non-consonant smile. Malocclusion was successfully treated with upper and lower fixed orthodontic appliance therapy. An aesthetic and stable occlusion was achieved at the completion of treatment. Fixed lingual and thermoplastic retainers were used to ensure stability of treatment outcome

**Key words:** Diastema, Malocclusion, Orthodontics

## Authors' Affiliation

<sup>a</sup>Department of Child Dental Health, Faculty of Dental Sciences, College of Medicine, University of Lagos, PMB 12003, Idiaraba, Lagos

<sup>b</sup>Department of Child Dental Health, Lagos University Teaching Hospital, PMB 12003, Idiaraba, Lagos.

## Correspondence:

Umeh Onyinye D  
Department of Child Dental Health,  
Faculty of Dental Sciences, College of Medicine, University of  
Lagos, PMB 12003, Idiaraba, Lagos  
Telephone number: +2348081412492  
E-mail: [umehod@gmail.com](mailto:umehod@gmail.com)

## Introduction

**M**idline diastema is characterized by the occurrence of space between the central incisors in the maxilla and less frequently in the mandible.<sup>1</sup>

The prevalence of diastema was found to be 75.3% (Caucasoid), 68.2% (Mongoloids), and 73.9%

(Negroids) in the maxilla, while for the mandible it was found to be 24.7% (Caucasoid), 31.8% (mongoloids), 26.1% (Negroids).<sup>2</sup> A Nigerian study found the incidence of midline diastema to be 26.1% (maxilla- 21.0%, mandible- 1.9%, and both arches- 3.2%),<sup>3</sup> while another found it to be 37%.<sup>4</sup> It has been shown that midline diastema is more common in the maxilla than in the mandible, with a slight female preponderance.<sup>3,4,5</sup>

Local factors like abnormal labial frenulum, missing maxillary lateral incisors, ectopic maxillary canines, missing teeth, anterior traumatic bite, tooth size - arch length discrepancies, mesio-distal angulation of the central incisors, inter-arch relationship, oral habits, presence of supernumerary tooth, pathologic lesions of the hard and soft tissue, iatrogenic causes, or physiologic development have been associated with the etiology of midline diastema. In addition to all these, genetic influences also play a vital role in its etiology.<sup>6,7</sup>

Midline diastema is usually considered a thing of beauty in this part of the world.<sup>5</sup> It was found that midline diastemas located in the maxilla, of about 2mm to 3mm in width, are preferred to their larger counterparts.<sup>8</sup> There has been reports of individuals going as far as artificially creating maxillary midline diastema.<sup>9</sup> The decision to treat is usually dependent on patient's preference and presence of any other associated occlusal discrepancies.<sup>5</sup>

The outcome of midline diastemas varies, some

resolving spontaneously while others may require intervention.<sup>10</sup> Possible therapeutic approaches to the management of midline diastemas requiring intervention may include either orthodontic, restorative, or surgical approach. Various combinations of the therapeutic approaches mentioned above may also be employed.<sup>11</sup>

### Diagnosis and Aetiology

A 23 year-old male with the chief complaint of large gaps between the upper and lower front teeth (maxillary and mandibular midline diastema). He was more concerned about the gap in the upper arch

proportions. The lips were mildly incompetent with a Jackson's lip pattern of 2 over 1, with a non-consonant smile (Fig 1). Intraoral evaluation revealed a full complement of teeth from the central incisors to the third molar in all quadrants. The molar relationships were Angle's Class I on both sides. Anterior overjet was 3mm and overbite was reduced and complete. The upper midline was coincident with the midline of the face, and a lower midline deviation to the right by 1 mm. There was also the presence of maxillary abnormal frenal attachment (Figure 1).

Examination of the upper arch revealed a moderate upper anterior segment spacing of 6.5 millimeters



**Figure 1: Pre-treatment facial and intraoral photographs**

and also wanted to close up other spaces between the teeth. He was eager to improve his appearance and self-esteem by undergoing orthodontic treatment. He was in good health with no significant systemic medical history. Extra oral evaluation revealed a concave profile with facial symmetry, good chin-throat support, soft tissue line, and normal vertical

inclusive of a midline diastema of 4 millimeters (mm), proclined upper incisors, and slight mesiolabial rotations of the upper left central incisors and canine (Figure 1). The lower arch revealed a 3mm midline diastema, 3mm posterior segment spacing, proclined incisors, rotation of the mandibular left canine, and an exaggerated curve of spee.

The panoramic radiograph showed the presence of all the teeth. Presence of periodontal compromise was evidenced by horizontal and vertical bone loss, especially on the mesial and distal root surfaces of the mandibular left second premolar and right second



**Figure 2: Pre-treatment panoramic radiograph**

molar, as well as the mesial surface of the mandibular left second molar (Figure 2).

Cephalometric analysis showed a Class I skeletal pattern 2 with increased upper and lower incisal angles, proclined upper and lower incisors and reduced interincisal angle (IIA), Frankfort mandibular plane angle (FMA) and lower facial



**Figure 3: Pre-treatment cephalometric radiograph**

height (LFH) (Figure 3, Table 1).

Based on these findings, the patient was diagnosed with Angles Class I malocclusion on Skeletal Pattern 2, complicated by maxillary and mandibular midline diastema, severe upper anterior segment spacing, mild lower posterior segment spacing. Also diagnosed were, proclined maxillary and mandibular incisors, low maxillary frenal attachment, lower midline deviation, mild rotations and exaggerated curve of spee in the mandible .

### Treatment Objectives

The orthodontic treatment objectives for this patient were to (1) close up the maxillary and mandibular midline diastema as well as other spaces (2) retrocline proclined upper and lower incisors (3) correct mandibular midline deviation (4) derotate rotated teeth (5) level the mandibular curve of spee (6) achieve lip competence and (7) achieve a consonant smile. These objectives were to be achieved while maintaining the stable occlusion, and the Class I molar relationship with the ultimate aim of improving the patient's appearance and self-confidence.

### Treatment Alternatives

The primary complaint of the patient was the midline diastema in the mandibular and maxillary arches. Based on literature review, the following treatment options were recommended:

1. Fixed appliance therapy, pre-adjusted edgewise; the non-extraction treatment. The options of using either the metal or ceramic Roth 022 slot and space closure on 019 X 025 stainless steel wire was suggested. This will be followed by frenectomy to reposition the frenal attachment at the end of space closure.
2. The use of aligners without extractions followed by the frenectomy procedure was the second option. This option is more aesthetic than the previous option. A major drawback to this option was the high cost of treatment relative to option 1.

The Retention plan was to use fixed lingual and thermoplastic retainers in the upper and lower arches.

After explaining the pros and cons of each alternative to the patient, he selected the first option for treatment.

### Treatment Progress

Treatment began with oral hygiene prophylaxis and



Figure 4: Intra treatment photographs



Figure 5: Post treatment facial and intraoral photographs

periodontal management of the periodontally compromised teeth. This was followed by bonding the teeth from the incisors to the second molars in all quadrants. Bondable buccal tubes were used on the molars instead of bands due to the presence of periodontal pockets around the molars at presentation. Levelling and alignment was achieved by sequential use of nickel titanium wires (014, 016, 018, 020 inches) and 020 inches stainless steel wires. The midline diastema and other intra arch diastema were closed by the use of elastic chain and active tie backs on 019 by 025 inches stainless steel wire. (Figure 4)

Active treatment took 19 months. Fixed lingual and thermoplastic retainers were used to maintain the positions of the teeth after treatment.

### Treatment Results

The teeth were well aligned and leveled at the end of treatment. The Class I molar and canine relationships were maintained. Lip competence was achieved with a satisfactory consonant smile (Figure 5)

Complete closure of the mandibular and maxillary midline diastema as well as other spaces within the

area was achieved (Figure 5). There was correction of the mandibular midline deviation, derotation of rotated teeth and leveling of the exaggerated curve of spee in the mandible.

The post treatment cephalometric analysis showed that the normal mandibular and maxillary incisal



**Figure 6: Post treatment cephalometric radiograph**

angles were achieved with satisfactory soft tissue balance. (Figure 6, Table 2).


Root parallelism was achieved and there was significant root resorption from the evaluation of the post treatment panoramic radiograph (Figure 7).

**Table 1: Pre-treatment cephalometric measurements**

Measurement	Pre-treatment values	Range for Nigerian norms	Comment
SNA	87.5°	85.5 ° ±3.5	Normal
SNB	82.5°	82.7 ° ±3.0	Normal
ANB	5°	2-4 °	Increased
U1 TO FP	128°	119-127 °	Increased
L1 TO MP	108°	96-104 °	Increased
IIA	101.5°	108-116 °	Reduced
FMA	23°	24-26 °	Reduced
LFH	53%	55%	Reduced

**Table 2: Post-treatment cephalometric measurements**

Measurement	Post-treatment values	Range for Nigerian norms	Comment
SNA	87.5°	85.5 ° ±3.5	Normal
SNB	83°	82.7 ° ±3.0	Normal
ANB	4.5°	2-4 °	Increased
U1 TO FP	124°	119-127 °	Normal
L1 TO MP	101°	96-104 °	Normal
IIA	112°	108-116 °	Normal
FMA	24.5°	24-26 °	Reduced
LFH	54%	55%	Reduced

**Figure 7: Post treatment panoramic radiograph****Figure 8: Pre and post treatment facial and intraoral photographs**

## Discussion

The presence of a midline diastema may pose both an aesthetic and psychological problem due to its prominent position in the dental arch, hence the constant request for correction by the patients, especially wide diastemas. Although multiple developmental, genetic and environmental factors have been incriminated in the aetiology of midline line diastema, it is believed to be mainly a multifactorial phenomenon.<sup>6,7</sup> Nonetheless, for effective management, accurate diagnosis and aetiology is required, putting into consideration also if it occurs as a single entity or in conjunction with other dental or occlusal anomalies.<sup>6,12,13</sup>

Management of midline diastema is usually interdisciplinary in approach; with treatment options ranging from simple or complex restorative treatment to orthodontic and surgical procedures depending on the aetiology. In this case report, the aetiology was a combination of tooth size-arch length discrepancy and high frenal attachment. Orthodontic treatment was considered due to the existence of other occlusal anomalies which were of concern to the patient, followed by frenectomy at the completion of orthodontic space closure.

In the orthodontic management of this patient, the use of bands as attachment on the molars were avoided to prevent further plaque accumulation, bacterial colonization and aggravation of the existing periodontal disease. Separator placement and band cementation have been associated with bacteremia.<sup>14</sup>

Treatment outcome in the case report was successful. The maxillary and mandibular midline diastema was closed at the end of treatment. Upper and lower spaces were closed and incisor proclination were corrected. Derotation of teeth was carried out and

overbite corrected. Lip competence and a consonant smile achieved. Most importantly, the periodontal health of the patient was preserved at the end of orthodontic treatment as seen in the post orthodontic treatment panoramic radiograph (Figure. 7).

The patient however declined the frenectomy procedure after completion of orthodontic treatment despite being informed extensively of the risk of relapse if not carried out. A fixed lingual retainer in conjunction with thermoplastic retainer was therefore employed as the retention protocol.

Retention protocol of midline diastema is of extreme importance for long term stability of treatment outcome, with aetiology of importance in this decision making.<sup>7</sup>

## Conclusion

Orthodontic treatment is a treatment option for the management of midline diastema. This case reports a case of maxillary and mandibular midline diastema occurring in conjunction with other occlusal anomalies. The aetiological factors implicated were tooth size- arch length discrepancy and abnormal frenal attachment. It was successfully treated using a fixed orthodontic appliance. The treatment outcome was satisfactory.

## Authors' Contributions

All the authors contributed to the conceptualization, design, and write-up of the manuscript.

## Funding/Grants

Self-funded

## Conflict of Interest

Nil

## References

1. Angle EH. Treatment of Malocclusion of the Teeth 7th edn. Philadelphia: SS White Dental Manufacturing Company, 1907; 103-104: 37-42
2. Lavelle CL. The distribution of diastemas in different human population samples. *European Journal of Oral Sciences*. 1970;78(1-4):530-534.
3. Omotosho GO, Kadir E. Midline diastema amongst South-Western Nigeria. *The Internet Journal of Dental Science*. 2010;8(2):27-32
4. Anosike AN, Sanu OO, Da Costa OO. Malocclusion and its impact on quality of life of school children in Nigeria. *West Afr Med J*. 2010;29(6):417-424.
5. da Costa OO. Midline diastema in a Northern Nigeria population. *Nig Qt. J Hosp Med*. 1996;6(4):287-289.
6. Abu-Hussein M, Watted N. Maxillary midline diastema—Aetiology and orthodontic treatment—clinical review. *IOSR J Dent Med Sci*. 2016;15(6):116-310.
7. Abrahams R, Kamath G. Midline diastema and its aetiology—a review. *Dental update*. 2014;41(5):457-464.
8. Akinboboye BD. Transcultural perception of maxillary midline diastema. *Int J Esthet Dent*. 2015;10:610-617.
9. Umanah A, Omogbai AA, Osagbemi B. Prevalence of artificially created maxillary midline diastema and its complications in a selected Nigerian population. *African health sciences*. 2015;15(1):226-232.
10. Huang WJ, Creath CJ. The midline diastema: a review of its etiology and treatment. *Pediatric dentistry*. 1995;17:171-179.
11. Gkantidis N, Kolokitha OE, Topouzelis N. Management of maxillary midline diastema with emphasis on etiology. *Journal of Clinical Pediatric Dentistry*. 2008;32(4):265-72.
12. Azzaldeen A, Muhamad AH. Diastema closure with direct composite: architectural gingival contouring. *Journal of Advanced Medical and Dental Sciences Research*. 2015 Jan;3(1):134-139.
13. Abu-Hussein M, Watted N, Abdulgani A. An Interdisciplinary Approach for Improved Esthetic Results in the Anterior Maxilla Diastema *Journal of Dental and Medical Sciences* 2015,14(12), 96-101
14. Umeh OD, Sanu OO, Utomi IL, Nwaokorie FO. Prevalence and intensity of bacteraemia following orthodontic procedures. *Int Orthod*. 2016;14(1):80-94.

# Instructions for Authors

West African Journal of Orthodontics is a peer-reviewed journal published by affiliated Orthodontic Groups and Associations in the West African Sub region. The journal gives priority to reports of outstanding clinical and experimental and epidemiological works on malocclusion, dento-facial defects as well as important contributions related to common orthodontic problems in children, adolescents and adults worldwide.

## Submission

Manuscripts and registered letters should be sent to: the Editor, West African Journal of Orthodontics, Department of Child Dental Health, Faculty of Dentistry, College of Health Sciences Obafemi Awolowo University, Ile-Ife, Osun State. Nigeria.

Manuscripts in MS word attachments may also be submitted via Email to [wajoeditorinchief@yahoo.com](mailto:wajoeditorinchief@yahoo.com), in addition to hard copies. Tables, figures and text should be included in the same file if possible. Authors may submit their research works by email only; such manuscripts need not be simultaneously sent by post.

However, photographs and/or figures need to be sent separately as hard copy (under figures and illustrations).

## Acceptance

Manuscripts should meet the following criteria: original material, clear writing, appropriate study methods, valid data, and reasonable conclusions supported by the data, in short, they should contain important information on topic of general orthodontic interest.

## Peer-review Process

All the manuscripts that adhere to its style and Instructions for Authors are referred to peer-review. Some of them are rejected immediately after an inhouse review. The rejection at this stage is due to insufficient originality, serious scientific flaws or absence of message. The remaining articles are sent to at least two reviewers who are experts in the subject. Manuscripts are reviewed with due respect for authors' confidentiality, and the identity of peer reviewers is also kept confidential. A decision is made from 6 to 12

weeks according to the response from reviewers, revision by the author(s) and reappraisal on the revision.

The accepted manuscripts are subjected to editorial revision to comply with the requirements on language and style of the journal. The rejected manuscript is not returned to authors but its copies are kept for 3 months to answer any queries. The copyright of the accepted and published articles is held by the journal and all the published materials cannot be reproduced or published elsewhere, in whole or part, without the written permission from the editor.

## Duplicate Submission

Manuscripts are considered with the understanding that they have not been published previously and are not under consideration by another publication. The author should alert the editor if the work includes subjects about which a previous report has been published. A research paper submitted to this journal should not overlap by more than 10% with the previously published material or work submitted elsewhere, which would be considered as duplicate publication. If in doubt, authors may forward copies of the published work or material submitted elsewhere to this journal for decision making.

## Proofs and Reprints

The corresponding author of the accepted article shall be supplied with the proof. Corrections on the proof should be restricted to errors only and no substantial additions/deletions should be made. No addition or deletion in the names of the authors is permissible at this stage. A copy of the issue carrying the article is supplied free of charge to the authors.

Reprints may be ordered on payment in advance.

## Categories of Articles

Articles can be sent as editorials, original articles, review articles, special communications, brief reports, case reports, letters to editor, commentaries, or for images section.

address. They are mostly included under Events of Interest free of cost. This journal reserves the right to be selective in publishing these announcements.

### **Preparing Manuscripts**

Manuscripts should be prepared in accordance with the Uniform Requirements for Manuscripts submitted to Biomedical Journals. 2 A summary of technical requirements for preparing the manuscript is provided below:

- Three copies of the manuscript should be submitted.
- Use 1 side of standard size 21.6x27.9 cm A4, white bond paper, with margins of at least 2.5 cm on each side.
- Double-space throughout including title page, abstract, text, acknowledgements, references, tables and figure legends. Start each of these sections (in same order) on a new page, numbered consecutively in the upper right hand corner, beginning with the title page.
- Use at least 12 point font size (Times New Roman or Arial).
- Submit photographs and transparencies in a separate heavy paper envelope (enclosed in cardboard, to prevent bending during mail handling).
- Conventional units are preferred with SI units in parenthesis, if available. The metric system is preferred for the expression of length, area, mass and volume.
- Use nonproprietary names of material rugs, devices and other products.
- All manuscripts should be accompanied by a signed statement by all authors regarding authorship, responsibility, financial disclosure and acknowledgements, as per standard format (Appendix J)[23 1 Those sending their manuscript through email are also required to submit this form by post with original signatures.

Manuscripts not fulfilling the technical requirements shall be returned to the authors without initiating the peer-review process.

### **Title Page**

The page should contain (i) the title of the article: which should be concise but informative (simpler the title the better; preferably it should contain all the key words to help electronic retrieval reliably); (ii) a short

running title of less than 40 characters placed at the foot end of the title page; (iii) initials and surname of each author with the highest academic degree(s) and designation at the time when the work was done; (iv) details of the contribution of each author; (v) name of department(s) and institution(s) to which the work should be attributed; (vi) disclaimers, if any; (vii) name, address, telephone, fax, email address of the corresponding author, (viii) source(s) of support in the form of grants, equipment, drugs or all of these; and (ix) declaration on competing interests.

### **Authorship**

All persons designated as authors should qualify for the authorship. Authorship credit should be based on substantial contributions to (i) concept and design, or acquisition of data, or analysis and interpretation of data; (ii) drafting the article or revising it critically for important intellectual content; and (iii) final approval of the version to be published. Conditions 1, 2 and 3 must all be met. Participation solely in the acquisition of funding or the collection of data does not justify authorship. All such people who contributed to the work but do not satisfy all the conditions should be listed in the acknowledgements.

Authors are responsible for obtaining written permissions from everyone acknowledged by name. One of the authors shall act as guarantor of the paper and he/she should take the responsibility for the integrity of the work as a whole, from its inception to published article.

Authors should provide a description of what each author contributed on the title page. Subsequently, no names can be added or deleted without written permission of the editor. Written consent of authors whose names are being deleted should be obtained.

This journal reserves the right to satisfy itself regarding the specific role of each listed author to justify authorship. All authors must give signed consent to publication (Appendix 1).

### **Competing Interest**

Competing interest for a given manuscript exists when the author has ties to activities that could inappropriately influence his or her judgment, whether or not judgment is in fact affected. Financial relationships with industry for example, through employment, consultancies, stock ownership, honoraria, expert testimony, either directly or through immediate family, are usually considered to be the most important competing interests. However, conflicts can

---

## Original Article

Original articles should report original research relevant to basic and clinical orthodontics including randomized trials, intervention studies, studies of screening and diagnostic tests, cohort studies, cost effectiveness analyses and case control studies. While reporting randomized controlled trials (RCT), authors must attempt to be in conformity with the consolidated standards of reporting trial.

## (CONSORT) statements

Each manuscript should be accompanied with a structured abstract (divided into background, methods, results and conclusions) in no more than 250 words. Four to five key words to facilitate indexing should be provided in alphabetical order along with the abstract. The text should be divided in sections on introduction, methods, results, discussion and conclusion.

Acknowledgment section may be included where necessary. Number of tables and figures should be limited to the very relevant ones and may be compressed if necessary. The typical text length for such contributions is 2500-3 500 words (excluding title page, abstract, tables, figures, acknowledgments and references).

## Brief Report

Short accounts of original studies are published as brief reports. The text should be divided into sections, i.e., abstract, introduction, methods, results and discussion.

Abstract should be of 100-150 words highlighting the aims, methods and main results along with 3-4 key words.

The text should contain no more than 1500 words, 3 illustrations or tables and up to 20 references, preferably recent publications.

## Review Article

State-of-the-art review articles or systematic, critical assessments of literature are also published. Normally a review article on a subject already published in the West African Journal of Orthodontics is not accepted for a period of 3 years.

The typical length for review articles is 2000-3000 words, excluding tables, figures, and references.

Authors submitting review manuscripts should include a structured abstract of around 200 words describing the need and purpose of review, methods used for selection, extraction and synthesis of data, and main conclusions.

Clinical cases highlighting uncommon malocclusion condition, orthodontic treatment techniques are published as case reports. Single case reports are usually not accepted, unless some new or unusual aspect regarding aetiopathogenesis, diagnosis or management is brought out that adds to the existing body of knowledge. The text should not exceed 1000 words and is divided into sections, i.e., abstract, introduction, case report and discussion. The number of tables/figures should be limited to 2. Ten recent references are acceptable. A maximum of 3 or 1 author is permitted from the principle and each of the associated departments respectively. Thus, case reports from only one investigative department can have a maximum of 3 authors.

## Letter to Editor(s)

Letters commenting upon a recent article in the West African Journal of Orthodontics are welcome.

Such letters should be received within 6 months of the article's publication. At the editorial board's discretion, a letter may be sent to authors! experts for comments and both letter and reply may be published together. Letters may also relate to other topics of interest to orthodontists and others, and/or useful clinical observations. Letters should not be more than 400 words. The number of authors should not exceed 2, including the authors' reply in response to a letter commenting upon an article published in this journal.

## Images Section

A short text of about 150 words depicting the condition with color photographs (vide infra) is needed.

Normally only clinical photographs are accepted but accompanying skiagrams or pathological images could also be considered for publication.

Photographs should be of high quality, clearly identify the condition and preferably add to the existing knowledge.

## Personal Viewpoint

Such articles are published on topical orthodontic issues including social aspects. It is expected that the authors have sufficient credible experience on the subject for giving viewpoints. These should not exceed 1500 words.

## Notes, News and Events of Interest

Announcements for conferences, symposia, meetings or courses may be sent for publication in advance. The announcements should provide title, date(s) and place of the event and contact address, telephone, and email

occur for other reasons, such as personal relationships, academic competition and intellectual passion. If any of the authors have accepted reimbursement for attending symposium, a fee for speaking, fee for organizing educational reach, funds for a member of the staff of consultation fees from an organization that may in: way gain or lose financially from the result of the study, review, editorial or letter, a competing interest would be deemed to exist. If any of the authors had been employed by an organization that may in any way gain or lose financially from the publication, or if any of them hold stocks or shares in such an organization, competing interest would be deemed to exist. If competing interest exists, the author(s) must disclose them while submitting the manuscript.

### **Abstract and Key Words**

The second page should carry an abstract in case of original article (250 words), review article (200 words), brief report (100-150 words), and case report (50 words), respectively. For original article and reviews, the abstract should be structured as detailed earlier. For brief reports, the abstract should state the purpose of the study, basic methodology, main findings (giving specific data and statistical significance) and key conclusion(s). Below the abstract, authors should provide 3-5 key words for indexing; terms from the Medical Subject Headings (MESH) list of Index Medicus should be used. The basic structure of a paper follows the well known acronym IMRAD, which stands for Introduction (what questions was asked), Methods (how was it studied), Results (what was found) and Discussion<sup>4</sup>.

### **Introduction**

The introduction must clearly state the question that the author(s) tried to answer in the study. It may be necessary to briefly review the relevant literature. Only cite those references that are essential to justify the proposed study.

### **Materials and Methods**

The methods section should describe, in a logical sequence, how the study was designed (e.g., how randomization was done), carried out (e.g., how subjects were chosen or excluded, ethical considerations, accurate details of materials used, exact drug dosage and form of treatment, etc.) and data were analyzed (e.g., an estimate of the power of the study, exact test used for statistical analysis, etc.). For standard methods, appropriate references are sufficient, but if standard methods are modified these should be clearly brought out.

Authors should provide complete details of any new methods or apparatus used (manufacturer's name and address in parentheses).

### **Ethics**

When reporting experiments on human subjects, authors should indicate whether the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional or regional) and with the Helsinki Declaration of 1964, as revised in 2000.

They should indicate whether the study was approved by the Institutions' Ethical Committee, and whether informed consent was obtained from the study participants. They should not use patients' names, initials, or hospital numbers, especially in illustrative material. This journal reserves the right to reject a manuscript on ethical grounds, on the basis of recommendations of its "Ethical Committee", even if the research has been cleared by the institutional ethical committee. Moreover, when reporting experiments on animals, authors should indicate whether the institutional and national guide for the care and use of laboratory animals was followed.

### **Statistics**

Authors should describe statistical methods with enough detail to enable a knowledgeable reader with access to the original data to verify the reported results. When possible, they meet to quantify findings and present them with appropriate indicators of measurement error or uncertainty (such as confidence intervals). Actual P values are provided rather than stating as just  $<0.05$  or  $>0.05$  etc. References for the design of the study and statistical methods should be to standard works when possible (with pages stated) rather than to papers in which the designs or methods were originally reported. Any general-use computer programs used should be specified and statistical terms, abbreviations, and most symbols be defined.

### **Results**

This section should include only relevant, representative data and not all information collected during the study. Major findings should be presented clearly and concisely. Text, tables, and illustrations should be used sensibly while avoiding repeating in the text all the data depicted in the tables or illustrations and emphasizing or summarizing only important observations. Tables and figures should be restricted to those needed to explain the argument of the paper and to assess its support. It is necessary to cite the tables in the text and type them on separate sheets. It may also be useful to mention what the study did not find.

## Discussion

Discussion ordinarily should not be more than one third of the total length of the manuscript. This section should include a summary of the major findings, their relationship to other similar studies, limitations of methods and implications of these findings in future research. Conclusions should be linked to the goals of the study. Unqualified statements and conclusions which are not completely supported by the data should be avoided. Authors should also refrain from making statements on economic benefits and costs unless their manuscript includes economic data and analyses.

## Acknowledgements

In acknowledgements section, it is suitable to list all contributors who do not meet the criteria for authorship, such as a person who provided purely technical help, writing assistance, or a department head who provided only general support. Financial and material support should also be acknowledged.

Groups of persons who have contributed materially to the paper but whose contributions do not justify authorship may be listed under a heading such as "clinical investigators" or "participating investigators", and their function or contribution should be described, for example, "served as scientific advisers", "critically reviewed the study proposal", "collected data", or "provided and cared for study patients". A written consent is required from all the persons acknowledged, indicating their acceptance for the same.

## Contributions to joint-authorship

In the case of multiple author-ship, authors are expected to state clearly their contributions to the paper being considered for publication in terms of study initiation, design including methodology, data collection, analysis and final write-up. The editorial board reserves the right to remove any author's name if the contribution is insignificant.

## References

References should be numbered consecutively in the order in which they are first mentioned in the text.

References are identified in text, tables, and legends by Arabic numerals in parentheses. References cited only in tables or in legends to figures should be numbered in accordance with the sequence established by the first identification in the text of the particular table or figure.

The titles of journals should be abbreviated according to the style used in Index Medicus. Authors are required not to use abstracts, unpublished observations and personal communications as references. References to papers accepted but not yet published should be designated as "in press"; authors should obtain written permission to cite such papers as well as verification that they have been accepted for publication.

The references must be verified by the author against the original documents. The Uniform Requirements style (the Vancouver style) is based largely on an American National Standards Institute (ANSI) standard style adapted by the NLM for its databases.

## Journal Article

List all authors when 6 or less. When 7 or more, list only first six and add et al. Ngan P, Yiu C, Hu A, Hagg U, Ei SHY, Gunel E. Cephalometric and occlusal changes following maxillary expansion and protraction. *Eur J Orthod* 1998; 20: 237-254.

## Organization as Author

Australian Dental Association Inc. An Australian Schedule of Dental Services and Glossary. 7th edn. Sydney: Australian Dental Association Inc., 1996.

## Complete Book

Department of Health. Shifting the balance of power within the NHS: securing delivery. London: Doll, 2001.

Clayton D, Hills M. Statistical models in epidemiology. Oxford: Oxford University Press, 1993.

Farkas LG. Anthropometry of the Head and Face, 2nd Edn, New York; Raven Press; 1994

Book Chapter Lekholm U, Zarb GA. Patient selection and preparation. In: Branemark P1, Zarb GA, Albrektsson T, editors.

Tissue integrated Prostheses: Osseointegration in Clinical Dentistry, Chicago: Quintessence; 1988,199-209

## Thesis and Dissertation

Yong SJ. Bone mineral density of normal Korean adults. Ph.D. Thesis. Seoul, Korea; 1989 Anozike, AN. Orthodontic treatment needs and its impact on oral health related quality of life in Lagos school children aged 12-16 years. FMCDs. Dissertation. Lagos, Nigeria; 2006

### Conference Proceedings

Marshall SJ, Rixon RC, Whiteford DN, Cumming JT. The OrthoForm 3-Dimensional Clinical Facial Imaging System. Proceedings of the 15th IFHE Congress 1998; 15:83-87.

### Dictionary and Similar References

Stedman's medical dictionary. 26th ed. Baltimore: Williams & Wilkins; 1995. Apraxia; p.11 9-120. Unpublished accepted material Leshner AI. Molecular mechanism of cocaine addiction. N Eng J Med. In Press 1996.

### Material from Internet

World Health Organization, 2002.  
www.who.int/mental-health/prevention/suicide (accessed August 1, 2004).

### Tables

Each table should be typed in double-space on a separate sheet of paper. Tables not submitted as photographs must be numbered consecutively (Arabic numerals) in the order of their first citation in the text, with a brief but self explanatory title for each.

Each column should have a short or abbreviated heading. Explanatory matters are placed in footnotes, not in the heading. In footnotes all nonstandard abbreviations that are used in each table should be explained adequately. Statistical measures of variations should be identified such as standard deviation and standard error of the mean. Be sure that each table is cited in the text. If data are used from another published or unpublished source, it is necessary to obtain permission and acknowledge them fully.

### Figures and Instructions

Figures should be professionally drawn and photographed; freehand or typewritten lettering is unacceptable. Instead of original drawings, X-ray films, and other material, sharp, glossy, black-and-white photographic prints of high quality are necessary, usually 127x 173 mm (5x7 in) but no larger than 203x254 mm (8x10 in) For color illustrations negatives or positive transparencies are provided, along with color prints. It is preferable to have the photograph in portrait form rather than in landscape form to fit easily into one column. Letters, numbers and symbols in photographs should be clearly legible.

Each figure should have a label pasted on its back indicating the number of the figure, author's name, and an arrow to mark the top and left side of the figure.

It is unacceptable to write on the back of figures or scratch or mark them by using paper clips, and to bend figures or mount them on cardboard. If photographs of individual/people are used, either the subjects must not be identifiable or their pictures must be accompanied by written permission to use the photograph. It is advisable to cover the eyes unless specifically need to be shown. If a figure has been published, the original source should be acknowledged and written permission from the copyright holder be obtained to reproduce the material. Figures should be numbered consecutively (Arabic numerals) according to the order in which they have been first cited in the text.

### Legends for Illustrations

Legends for illustrations should be typed or printed out in double-space, starting on a separate page, with Arabic numerals corresponding to the illustrations.

When symbols, arrows, numbers, or letters are used to identify parts of the illustrations, each of them must be identified and explained in the legend. The internal scale should be explained and the method of staining in photomicrographs be identified.

### Units of Measurement

Measurements of length, height, weight, and volume should be reported in metric units, i.e., meter(m), gram(g), or liter(l) or their decimal multiples.

Milliliter or deciliter should be expressed as ml or dl.

Red and white blood cell counts are to be expressed as  $63 \times 10^6 / \text{mc l}$  and  $\times 10^6 / \text{mc}$  respectively. Temperatures should be given in degrees Celsius and blood pressures in millimeters of mercury (mmHg). All hematological and clinical chemistry measurements should be reported in the conventional system or in terms of the International System of Units (SI).

### Abbreviations and symbols

Only standard abbreviations are used in the text while avoiding abbreviations in the title and abstract.

The full term for which an abbreviation stands should precede its first use in the text unless it is a standard unit of measurement. Year, month, day, hour, minute and second should be abbreviated as yr, mon, d, h, mm, and s in tables respectively.

## References

1. Mother M, Schulz KF, Altman DG, for the CONSORT Group. The CONSORT statement Revised recommendations for improving the quality of reports of parallel group randomize Trials. *Lancet* 2001; 357: 1191-1194. (Also available from: URL: <http://www.consort-statement.org/>). Accessed June 28, 2002.
2. International Committee of Medical Journal Editors. Uniform Requirements for Manuscripts Submitted to Biomedical Journals. *Ann Intern Med* 1997;126:36-47. (Updated October 2001 version Available from: URL: <http://www.icmje.org/>). Accessed June 28,2002.
3. JAMA Instructions for Authors. Available from URL: <http://jama.ama-assn.org/>. Accessed June 28, 2002.
4. Hall GM. Structure of a scientific paper. In: Hall GM, ds. *How to write a paper*. London:BMJ Books, 2000.
5. 52nd WMA General Assembly. World Medical Association Declaration of Helsinki. Ethical principles for medical research involving human subjects. Available from: URL: <http://www.wma.net/>. Accessed June 28,2002.

## Appendix 1:

### Declaration of Originality and Transfer of Copyright

(Please download from Nigerian Association of Orthodontists (NAO) website <https://www.nao-ng.org/>)

This form is to be submitted with the initial copies of the manuscript to: West African Journal of Orthodontics, Department of Child Dental Health, Obafemi Awolowo University Ile-Ife, Osun State. Nigeria Manuscript No. (If known):

The author(s) hereby affirms that the submitted manuscript entitled:

I/We certify that the manuscript represents valid work and that neither this manuscript nor one with substantially similar content under my/our authorship has been published or is being considered for publication elsewhere. For papers with more than I author, we agree to allow the corresponding author to serve as the primary correspondent with the editorial office, to review the edited typescript and proof.

I/We have seen and approved the submitted manuscript. All of us have participated sufficiently in the work to take public responsibility for the contents. All the authors have made substantial contributions to the intellectual content of the paper and fulfill at least 1 condition for each of the 3 categories of contributions: i.e., Category 1 (conception and design, acquisition of data, analysis and interpretation of data), Category 2 (drafting of the manuscript, critical revision of the manuscript for important intellectual content) and Category 3 (final approval of the version to be published).

I/We also certify that all my/our affiliations with or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript are completely disclosed on the title page of the manuscript. My/our right to examine, analyze, and publish the data is not infringed upon by any contractual agreement.

I/We certify that all persons who have made substantial contributions to the work reported in this manuscript (e.g., data collection, writing or editing assistance) but who do not fulfill the authorship criteria are named along with their specific contributions in an acknowledgment section in the manuscript. If an acknowledgment section is not included, no other persons have made substantial contributions to this manuscript.

I/We also certify that all persons named in the acknowledgment section have provided written permission to be named.

The author(s) undersigned hereby transfer(s), assign(s), or otherwise convey(s) all copyright ownership, including any and all rights incidental thereto, exclusively to the West African Journal of Orthodontics, in the event that such work is published in the West African Journal of Orthodontics.

Authors name(s) in order of appearance in the manuscript; signatures (date):

