

WEST AFRICAN JOURNAL OF ORTHODONTICS

VOLUME 14, NUMBER 1

Print ISSN: 2315-9634

E-ISSN: 3141-5822

June 2025

Malocclusion, fingerprints and blood group

Cephalometric measurements and Photogrammetry

Pattern of malocclusion seen at AKTH

Artificial Intelligence in Orthodontics

Talon Cusps: Conservative management



Editor-in-Chief and Editorial Office

Prof. O.D Otuyemi (Nigeria)
Department of Child Dental Health,
Obafemi Awolowo University, Ile-Ife,
Osun State, Nigeria.

Deputy Editor in-Chief

Dr. I.G Isiekwe (Nigeria)
Department of Child Dental Health,
Faculty of Dental Sciences,
College of Medicine, University of Lagos,
Lagos, Nigeria.

Associate Editors

Prof. P.I Ngom (Senegal)
Prof. M. Newman-Nartey (Ghana)
Dr. G.K Amoah (Ghana)
Dr. T.A Yemitan (Nigeria)
Prof. O.O daCosta (Nigeria)
Dr. T.E Adeyemi (Nigeria)
Dr. A. Traore-Shumbusho (Nigeria)
Prof. O.O Sanu (Nigeria)
Dr. M.A Ernest (Nigeria)
Dr. O. D Umeh (Nigeria)
Dr. O. A Aghimien (Nigeria)

Advisory Editorial Board

Prof. A. Harris (South Africa)
Dr. H.K Chuan (Malaysia)
Dr. D. Mahony (Australia)
Dr. H. Bellardie (UK)
Dr. J. Noar (UK)
Prof. P. Hlongwa (South Africa)
Prof. M.C Isiekwe (Nigeria)
Dr. A. Soyombo (Nigeria)
Dr. R. Chamda (South Africa)
Dr. D. Fashemo (USA)

Website: www.nao-ng.com

Contents

Editorial	3
------------------	---

Sanu O.O.

Original Articles

The Relationship Between Skeletal Malocclusions, Fingerprints and Blood Group in Patients Attending an Orthodontic Clinic in Benin City	4
--	---

Osemwegie C, Umweni AA, Omoregie F.O

Correlations between Cephalometric Measurements of Hard Tissues and Photogrammetric Features of Facial Soft Tissues	20
--	----

Djibril Cisse, Alpha Badiane, Malick Guèye, Irado Rakoto Solofo, Joseph Samba Diouf, Khady Diop-Bâ, Papa Ibrahima Ngom

Pattern of Malocclusion seen at Aminu Kano Teaching Hospital, Kano Nigeria: A 5 Year Review	29
--	----

Jibril M, Oguchi C, Adeyemi TE, Yahaya A.

The Role of Artificial Intelligence in Orthodontics	40
--	----

Awwal NM, Aladejare AL

Talon Cusp: An Uncommon Occlusal anomaly – Conservative Management of two Cases	50
--	----

Agboghoroma GO, Akinyemi OA, Enabulele JE.

Instructions for Authors	56
---------------------------------	----

Editorial

Malocclusion significantly affects individuals' quality of life, impacting nutrition, speech, and psychosocial well-being. In low- and middle-income countries (LMICs), access to orthodontic care is limited—not because of a lack of need, but due to the chronic shortage of trained specialists. At the heart of this problem lies a fragile training infrastructure plagued by underfunding, inconsistent standards, and the relentless loss of talent to more developed nations.

Dental schools in many LMICs operate under tight financial constraints, often relying on outdated equipment and lacking essential clinical tools like imaging systems or modern orthodontic materials. Students graduate with minimal exposure to contemporary practices, which undermines their confidence and readiness for clinical work. Meanwhile, the lack of simulation technologies and hands-on models limits the development of essential skills during training.

This is further complicated by restricted internet access and the high cost of digital resources, which keep many faculty and students cut off from global developments in orthodontics. In an age where digital tools drive education and research, this technological divide deepens inequality and slows the pace of local progress. Then there's the ongoing loss of human capital. Orthodontists trained in LMICs often leave in search of better working conditions, research opportunities, and financial security abroad. This migration is a major setback, draining already limited health systems of expertise, leadership, and mentorship. In effect, LMICs are subsidizing the specialist workforce of wealthier nations.

Curriculum design also poses challenges. In many institutions, orthodontic training includes excessive rotations outside of the specialty and lacks emphasis on critical thinking, research, and innovation. This creates a generation of clinicians who are technically competent but disconnected from the evidence base and unprepared to contribute to the evolution of care that meets local needs.

To move forward, orthodontic education in LMICs must become more intentional and adaptive. Curricula should be aligned with population needs, focusing on high-impact clinical skills, ethical care, and region-specific challenges. Affordable simulation tools and open-access digital learning can help bridge resource gaps. Equally important is faculty development—training the trainers—to foster leadership, mentorship, and academic growth within local institutions.

National dental associations must also play a more active role in setting training standards, supporting quality assurance, and advocating for investment in dental education. Governments, in turn, must create an enabling environment for professionals to stay and thrive, through better infrastructure, fair compensation, and research funding. International partners can support these efforts by promoting equitable collaboration and respecting ethical recruitment guidelines.

Improving orthodontic training in LMICs is not a luxury—it is a necessity. These countries deserve self-sustaining systems that produce skilled, motivated professionals equipped to meet the needs of their communities. With targeted investment, strong leadership, and strategic partnerships, that future is within reach.

Prof O.O. Sanu
Associate Editor

References

1. Bataringaya A, Ssenyonga R, Macgregor R. Challenges of setting up an orthodontic service in a developing country: Uganda. *Am J Orthod Dentofacial Orthop*. 2005;127:367–369.
2. Normando D. Challenges for orthodontic education in developing countries. *Dental Press J Orthod*. 2020;25:11–12.
3. Al-Sanabani JS, et al. Virtual reality in orthodontic education: a systematic review. *B3..MC Med Educ*. 2021;21:1–11.
4. World Health Organization. Global Code of Practice on the International Recruitment of Health Personnel. WHO; 2010.

The Relationship Between Skeletal Malocclusions, Fingerprints and Blood Group in Patients Attending an Orthodontic Clinic in Benin City

Osemwegie C,^a Umweni AA,^{a,b} Omoregie F.O^{c,d}

Abstract

Background: Documented evidence suggests a relationship between dermatoglyphic patterns, blood group system, and the development of malocclusion. The wide variations observed have been suggested to be due to genetic and environmental influences. More research to ascertain the relationship between malocclusion with fingerprints and blood group in Nigerians is essential. The objective of the study was to assess the relationship between skeletal malocclusion with fingerprints and blood group in a Nigerian population.

Methods: A total of 205 patients, aged 13-40 years, attending the Orthodontic clinic of the University of Benin Teaching Hospital, Benin City who met the inclusion criteria were selected. Their sagittal and vertical jaw relationships were determined by tracing the patients' lateral cephalometric radiograph. Fingerprints of both hands were taken using a Bio scanner, while blood samples were collected to determine their ABO blood group.

Results: More than half of the patients (55.6%) had skeletal pattern 1 malocclusion. The fingerprint pattern showed that the ulnar loop had the highest frequency across all types of skeletal malocclusion (61.8%), followed by whorl (24.5%), arch (10.2%), and radial loop (3.5 %). The fingerprint patterns of the left thumb, index finger, fifth finger, and right thumb had significant relationship with the different classes of skeletal malocclusion ($p < 0.05$)

Conclusion: There was a relationship between dermatoglyphic pattern and skeletal malocclusion. No significant relationship was found between skeletal malocclusion and blood group. Dermatoglyphic patterns of some digits show predictive values when compared to others.

Key words: Fingerprint, blood group, malocclusion, skeletal.

Authors' affiliations

^aDepartment of Orthodontics, University of Benin Teaching Hospital, Benin City, Edo State.

^bDepartment of Orthodontics, University of Benin, Benin City, Edo State.

^cDepartment of Oral Pathology and Oral Medicine, University of Benin, Benin City, Edo State.

^dDepartment of Oral Pathology and Oral Medicine, University of Benin Teaching Hospital, Benin City, Edo State.

Correspondence:

Osemwegie Collins
Department of Orthodontics
University of Benin,
Benin City, Edo State
Telephone Number: +2348032431452
Email Address: collinopeperano@yahoo.com
ORCID: 0009-0005-9011-0194

Introduction

Malocclusion may be defined as a significant deviation from what is defined as normal or ideal occlusion.¹⁻³ It is one of the common oral conditions that is often reported in

literature.¹ The aetiology of malocclusion is multifactorial, ranging from genetic factors to environmental factors and these factors influence the development of the jaws.⁴ Skeletal malocclusion occurs as a result of the distortion of the normal maxillary and/or mandibular growth during embryonic development.⁴ At present, malocclusions are ranked third in the order of priorities among the problems of dental public health worldwide.⁵ It is estimated that approximately 1.2 million people in the United States have an extreme dentoskeletal anomaly.⁶ Approximately 60% have a class II malocclusion, 25% have a class III malocclusion, and around 15% have a mix of dentoskeletal irregularities.⁷ Class III malocclusion is thought to be one of the most complex as well as challenging orthodontic problems to deal with. The prevalence of class III malocclusion in North Indian population is up to 3.4%, while the prevalence of class III malocclusion in Caucasians varies from 0.8 to 4.0% and rises to 13% in Japanese and Chinese populaces.⁷

Skeletal malocclusion is found to show characteristic dermatoglyphic patterns as explained by embryological origin of oral and dermatoglyphic patterns.⁸ Facial development begins as early as the 4th week of gestation.^{1,8} Palate begins to develop in the 6th week and is completed by the 12th week of gestation. The digital ridge patterns are indicators of growth-related developmental patterns, which start to appear during the 12th week of gestation, to be completed by 24th week, after which they remain throughout life.⁹

A relationship has also been found to exist between the ABO blood group system and some diseases and maxillofacial deformities.¹⁰⁻¹² The ABO blood group is genetically determined as they are inherited co-dominantly through genes on chromosome 9 from both parents.⁴ A person's ABO type is determined by the inheritance of 1 of 3 alleles (A, B, or O).¹⁰ The presence or absence of these antigens results in the four blood groups: A, B, AB, and O.⁹

Malocclusion leads to long-term complications such as temporomandibular joint (TMJ) dysfunctions, periodontal disease, obstructive sleep apnea, psychological disorders, and articulation errors.¹³⁻¹⁵

Late diagnosis of skeletal malocclusions leads patients to orthognathic surgery,¹⁶ therefore, the orthodontist must have sound knowledge of dental occlusion and the underlying skeletal relationship of the patient to arrive at the correct diagnosis and treatment plan for the malocclusion.⁵

Dermatoglyphic investigation is a method of exploring the genetic associations of malocclusion¹⁷ due to the fact that there is a strong association between dermatoglyphic patterns and sagittal skeletal discrepancies.¹⁸ A study by Reddy et al.¹⁹ found a significant association between different malocclusions and specific types of ridge pattern.

The relationship between the ABO blood group system and some oral diseases has been one of the most important human genetic traits.²⁰ A relationship exists between malocclusion and ABO blood group system, and given that both are related to genetic components, it may be stated that blood groups have an association with malocclusions.^{4, 10} The strongest

association between ABO blood group system and malocclusion was reported to be highest in blood group B, and least in blood group AB.⁴ Dermatoglyphic and ABO blood group studies done in association with malocclusion yielded diverse results, which could be due to the fact that numerous other factors such as ethnic and racial variations, congenital, environmental, and other local factors can also influence the development of malocclusions.^{19,21-23} More research is essential to ascertain the relationship between malocclusion with fingerprints and the blood groups in the Nigerian population.

This study is relevant to the field of orthodontics, particularly in preventive and interceptive orthodontics, for early diagnosis and correction of deviated growth patterns of the jaws. The objective of this study is to assess the relationship between fingerprint pattern and blood group with skeletal malocclusion in Nigerians.

Materials and Methods

Study design: This was a prospective cross-sectional study.

Study population and location: Patients seeking orthodontic treatment between the ages of 13 to 40 years old who met the selection criteria. The study was carried out in the Orthodontic clinic of the Department of Orthodontics, University of Benin Teaching Hospital, Benin City.

Selection criteria:

Inclusion criteria:

1. Patients with no history of orthodontic treatment.
2. Patients aged 13 years to 40 years old.
3. All patients who attended the orthodontic clinic, and gave written informed consent to participate in the study.
4. Patients without periodontal disease that may affect the positioning of the teeth.
5. Patients with full complement of teeth, excluding third molars.
6. All subjects were of Nigerian descent.

Exclusion criteria:

1. Patients with malformation syndromes or other craniofacial abnormalities associated with maxilla and mandible, and those with facial asymmetry and acquired skeletal defects.
2. Patients with congenital or acquired deformities of the fingers and amputated fingers.
3. Patients with skin diseases, with wounds or scars on the fingers.
4. Patients with periodontal conditions that may affect the positioning of the teeth.
5. Patients with previous extraction of the teeth, particularly the first molar.

Sample size determination and selection

The sample size was calculated using quantitative data formula:

$$n = \left[z \frac{\alpha}{2} \frac{\sigma}{E} \right]^2$$

$$= [13.62]^2 = 185.55 = 186$$

Where n is the sample size, a 95% degree confidence interval corresponds to $\alpha=0.05$, z the standard normal deviation, E the margin of error $E=1$ and standard deviation=6.95.

Materials and methods

The sagittal jaw relation was determined from the patient's lateral cephalogram with assessment of the following parameters: SNA, SNB, ANB and FMA.¹⁸ Cephalometric norms of the Nigerian population were used to group the patients' skeletal patterns (appendix I).²⁴ The type of malocclusion was classified using Steiner's analysis for skeletal malocclusion (ANB angle) into three groups: Skeletal Class 1, Skeletal Class 2 and Skeletal Class 3.^{16,25} The FMA was used to determine the vertical jaw relationship.²⁶

The fingerprints recording was done by the researcher, the hands of the subjects were cleaned by the researcher with soap and water and dried with sterile gauze. The impression of individual digits was

made by placing the bulb of the digits onto the surface of the bioscanner (see appendix IIa). The fingerprints were recorded with IBS software and analysed using the classification by Galton in 1892^{1,23} as shown in Appendix IIIb.

The blood test of the individual was done to determine their blood group by the researcher, in conjunction with a medical laboratory scientist.²⁷ Antisera A, B, and D were used, with blood samples from the participants (appendix IV). The presence of agglutination with antiserum A confirmed blood group A. The presence of agglutination with antiserum B confirmed blood group B. The absence of agglutination on both antisera A and B confirmed blood group O, and the presence of agglutination on both antisera A and B confirmed blood group AB. The presence of agglutination with antiserum D confirmed Rhesus positive blood group. The absence of agglutination with antiserum D confirmed Rhesus negative blood group. The patient's blood group was determined and recorded on a data collection form (see appendix V).

Ethical considerations

Ethical approval for this study was obtained from the Ethics and Research Committee, University of Benin Teaching Hospital, Benin City, Edo State, Nigeria (protocol number: ADM/E22/A/VOL. VII/14682). Informed consent was obtained from study participants or their parents/ caregiver for minors. Each participant filled the informed consent form and the minor gave assent before they were recruited into the study.

Results**Sociodemographic characteristics**

A total of 205 participants were examined, among them, 121 (59.0%) were females and 84 (41.0%) were males. The mean age was 24.86 ± 7.02 and the age ranged from 13 – 40 years, and the most common age group was 21 – 25 years (28.8%). A majority of the patients had blood group O, 143 (69.8%), the ulnar loop was the most common dermatoglyphic pattern at 1266 (61.8%), while skeletal class 1 at 114 (55.6%) was the most common skeletal malocclusion.

Table 1a: Distribution of patients' characteristics

Characteristic	Frequency N = 205	(%)
Age group (in years)		
13 – 15	16	7.8
16 – 20	46	22.4
21 – 25	59	28.8
26 – 30	41	20.0
31 – 35	19	9.3
36 – 40	24	11.7
Gender		
Male	84	41.0
Female	121	59.0
Blood group		
A	29	14.1
B	24	11.7
AB	9	4.4
O	143	69.8
Rhesus 'D' factor		
Positive	187	91.2
Negative	18	8.8
Dermatoglyphic pattern (¹N=2,050)		
Arch	210	10.2
Radial loop	71	3.5
Ulnar loop	1266	61.8
Whorl	503	24.5
Skeletal malocclusion pattern		
1	114	55.6
2	34	16.6
3	57	27.8

Table 1b: Vertical and sagittal skeletal measurement in degrees

ANB (Median (IQR))	3.0 (1.0 – 4.0)
FMA (Median (IQR))	22.0 (20.5 – 24.0)

Table 2: Association of dermatoglyphic patterns with skeletal malocclusion.

	Fingerprint pattern	Skeletal pattern 1 (%)	Skeletal pattern 2 (%)	Skeletal pattern 3 (%)	p-value
Left fifth finger	Arch	6 (5.3)	0 (0)	0 (0)	<0.001
	radial loop	2 (1.8)	0 (0)	0 (0)	
	ulnar loop	92 (80.7)	26 (76.5)	57 (100)	
	Whorl	14 (12.3)	8 (23.5)	0 (0)	
Left fourth finger	Arch	6 (5.3)	2 (5.9)	6 (10.5)	0.781
	radial loop	6 (5.3)	0 (0)	2 (3.5)	
	ulnar loop	73 (64.0)	24 (70.6)	36 (63.2)	
	Whorl	29 (25.4)	8 (23.5)	13 (22.8)	
Left index finger	Arch	20 (17.5)	1 (2.9)	11 (19.3)	0.0345
	radial loop	8 (7.0)	8 (23.5)	3 (5.3)	
	ulnar loop	45 (39.5)	16 (47.1)	25 (43.9)	
	Whorl	41 (36.0)	9 (26.5)	18 (31.6)	
Left middle finger	Arch	19 (16.7)	9 (26.5)	8 (14.0)	0.603
	radial loop	4 (3.5)	0 (0)	0 (0)	
	ulnar loop	68 (59.6)	19 (55.9)	38 (66.7)	
	Whorl	23 (20.2)	6 (17.6)	11 (19.3)	
Left thumb	Arch	20 (17.5)	5 (14.7)	8 (14.0)	0.006
	radial loop	3 (2.6)	5 (14.7)	0 (0)	
	ulnar loop	55 (48.2)	10 (29.4)	19 (33.3)	
	Whorl	36 (31.6)	14 (41.2)	30 (52.6)	
Right fifth finger	Arch	5 (4.4)	0 (0)	2 (3.5)	0.534
	radial loop	2 (1.8)	0 (0)	0 (0)	
	ulnar loop	95 (83.3)	32 (94.1)	53 (93.0)	
	Whorl	12 (10.5)	2 (5.9)	2 (3.5)	
Right fourth finger	Arch	3 (2.6)	3 (8.8)	2 (3.5)	0.207
	radial loop	0 (0)	0 (0)	2 (3.5)	
	ulnar loop	75 (65.8)	23 (67.6)	40 (70.2)	
	Whorl	36 (31.6)	8 (23.5)	13 (22.8)	
Right index finger	Arch	15 (13.2)	5 (14.7)	8 (14.0)	0.683
	radial loop	9 (7.9)	4 (11.8)	5 (8.8)	
	ulnar loop	61 (53.5)	15 (44.1)	23 (40.4)	
	Whorl	29 (25.4)	10 (29.4)	21 (36.8)	
Right middle finger	Arch	11 (9.6)	7 (20.6)	5 (8.8)	0.302
	radial loop	0 (0)	0 (0)	1 (1.8)	
	ulnar loop	87 (76.3)	25 (73.5)	43 (75.4)	
	Whorl	16 (14.0)	2 (5.9)	8 (14.0)	
Right thumb	Arch	14 (12.3)	6 (17.6)	3 (5.3)	0.048
	radial loop	7 (6.2)	0 (0)	0 (0)	
	ulnar loop	51 (44.7)	10 (29.4)	30 (52.6)	
	Whorl	42 (36.8)	18 (52.9)	24 (42.1)	

Fisher's exact test

The association between dermatoglyphic patterns and skeletal malocclusion is shown in Table 2. Arch was the most seen in the left thumb, index, and middle fingers of individuals with skeletal pattern I malocclusion. It was absent in the left fifth fingers of participants with skeletal patterns II and III malocclusion. Radial loop was most seen in the left index finger and right index finger of individuals with skeletal pattern II malocclusion. However, it was absent in the left fifth finger, left fourth finger, left middle finger and right fifth finger of participants with skeletal pattern II malocclusion. Ulnar loop pattern was most seen in the right and left fifth fingers of participants with skeletal pattern I malocclusion. Whorl pattern was seen more in the right thumb of individuals with skeletal

pattern III and left index finger of individuals with skeletal pattern I malocclusion. But the whorl pattern was absent in the left fifth finger and right middle finger of participants with skeletal pattern III malocclusion. A significant association exists between dermatoglyphic patterns and skeletal malocclusion, with the ulnar loop patterns of left fifth finger ($p = <0.001$), in association with skeletal pattern III and left index finger ($p = 0.035$) in associated with skeletal pattern II malocclusion. The whorl pattern of the left thumb ($p = 0.006$) is associated with skeletal pattern III and the whorl pattern of the right thumb ($p = 0.048$) was found to be associated with skeletal pattern II malocclusion

Table 3: Correlation of dermatoglyphic patterns of study participants with categories of Skeletal malocclusion

Digit	Dermatoglyphic pattern	Skeletal pattern 1 <i>N=114</i> No(%)	Skeletal pattern 2 <i>N=34</i> No(%)	Skeletal pattern 3 <i>N=57</i> No(%)	Kendall's tau (τ_b)	p-value
Left Thumb	Arch	20 (17.5)	5 (14.7)	8 (14.0)	0.136	0.029
	radial loop	3 (2.6)	5 (14.7)	0 (0)		
	ulnar loop	55 (48.2)	10 (29.4)	19 (33.3)		
	Whorl	36 (31.6)	14 (41.2)	30 (52.6)		
Left Index finger	Arch	20 (17.5)	1 (2.9)	11 (19.3)	-0.030	0.636
	radial loop	8 (7.0)	8 (23.5)	3 (5.3)		
	ulnar loop	45 (39.5)	16 (47.1)	25 (43.9)		
	Whorl	41 (36.0)	9 (26.5)	18 (31.6)		
Left Middle finger	Arch	19 (16.7)	9 (26.5)	8 (14.0)	0.013	0.835
	radial loop	4 (3.5)	0 (0)	0 (0)		
	ulnar loop	68 (59.6)	19 (55.9)	38 (66.7)		
	Whorl	23 (20.2)	6 (17.6)	11 (19.3)		
Left Ring finger	Arch	6 (5.3)	2 (5.9)	6 (10.5)	-0.037	0.578
	radial loop	6 (5.3)	0 (0)	2 (3.5)		
	ulnar loop	73 (64.0)	24 (70.6)	36 (63.2)		
	Whorl	29 (25.4)	8 (23.5)	13 (22.8)		
Left Fifth finger	Arch	6 (5.3)	0 (0)	0 (0)	-0.019	0.727
	radial loop	2 (1.8)	0 (0)	0 (0)		
	ulnar loop	92 (80.7)	26 (76.5)	57 (100)		
	Whorl	14 (12.3)	8 (23.5)	0 (0)		
Right Thumb	Arch	14 (12.3)	6 (17.6)	3 (5.3)	0.097	0.104
	radial loop	7 (100.0)	0 (0)	0 (0)		
	ulnar loop	51 (44.7)	10 (29.4)	30 (52.6)		
	Whorl	42 (36.8)	18 (52.9)	24 (42.1)		
Right Index finger	Arch	15 (13.2)	5 (14.7)	8 (14.0)	0.049	0.447
	radial loop	9 (7.9)	4 (11.8)	5 (8.8)		
	ulnar loop	61 (53.5)	15 (44.1)	23 (40.4)		
	Whorl	29 (25.4)	10 (29.4)	21 (36.8)		

Right Middle finger	Arch	11 (9.6)	7 (20.6)	5 (8.8)	-0.037	0.569
	radial loop	0 (0)	0 (0)	1 (1.8)		
	ulnar loop	87 (76.3)	25 (73.5)	43 (75.4)		
	Whorl	16 (14.0)	2 (5.9)	8 (14.0)		
Right Ring finger	Arch	3 (2.6)	3 (8.8)	2 (3.5)	-0.108	0.090
	radial loop	0 (0)	0 (0)	2 (3.5)		
	ulnar loop	75 (65.8)	23 (67.6)	40 (70.2)		
	Whorl	36 (31.6)	8 (23.5)	13 (22.8)		
Right Fifth finger	Arch	5 (4.4)	0 (0)	2 (3.5)	-0.046	0.470
	radial loop	2 (1.8)	0 (0)	0 (0)		
	ulnar loop	95 (83.3)	32 (94.1)	53 (93.0)		
	Whorl	12 (10.5)	2 (5.9)	2 (3.5)		

Table 3 shows the distribution and relationships of the categories of skeletal malocclusion to the participants' fingerprint patterns. When considered in categories, the proportion of participants with the whorl fingerprint pattern on the left thumb appeared to increase as the class of skeletal malocclusion increased. This monotonic relationship was confirmed on Kendall's tau-b test, which demonstrated a weak positive correlation of fingerprint patterns of the left thumb with increasing classes of skeletal malocclusion, which was statistically significant ($\tau_b = 0.136$, $p = 0.029$). The ulnar loop pattern was the dominant fingerprint

pattern in the left index and fifth finger, even though it did not appear to demonstrate a clear monotonic relationship with the classes of skeletal malocclusion as in the thumb. The fingerprint patterns on the other digits of the left hand were not related to the classes of skeletal malocclusion in the study participants.

On the right hand, all participants with a radial loop on their thumbs had the least class of skeletal malocclusion (100% vs 0%, $p = 0.043$). No clear monotonic relationship was found between participants' fingerprint patterns on any other digit and the class of skeletal malocclusion ($p > 0.05$).

Table 4a: Relationship between participants' dermatoglyphic patterns and quantitative measures of skeletal malocclusion

Hand	Digit	Dermatoglyphic pattern	No(%)	ANB (°) Median (IQR)	p-value	FMA (°) Median (IQR) [†]	p-value
Left	Thumb	Arch	33 (16.1)	3.0 (2.0 – 4.0)	0.127	22.0 (21.1 – 24.8)	0.141
		Radial loop	8 (3.9)	6.0 (2.0 – 9.0)		24.0 (21.3 – 28.3)	
		Ulnar loop	84 (41.0)	2.8 (2.0 – 4.0)		22.0 (20.6 – 23.0)	
		Whorl	80 (39.0)	3.0 (1.0 – 4.0)		21.8 (20.0 – 24.0)	
	Index finger	Arch	32 (15.6)	2.0 (-0.5 – 3.0)	0.037^a	22.0 (20.0 – 23.9)	0.029^b
		Radial loop	19 (9.3)	3.5 (2.0 – 5.0)		23.0 (22.0 – 25.0)	
		Ulnar loop	86 (42.0)	2.5 (1.3 – 4.0)		21.5 (20.0 – 23.0)	
		Whorl	65 (31.7)	3.0 (1.0 – 4.0)		22.0 (20.5 – 24.0)	

Middle finger	Arch	36 (17.5)	3.0 (2.0 – 5.0)	0.723	22.5 (22.0 – 25.0)	0.026^c
	Radial loop	4 (2.0)	2.5 (2.0 – 3.0)		21.0 (20.1 – 21.9)	
	Ulnar loop	125 (61.0)	3.0 (1.0 – 4.0)		22.0 (20.0 – 23.0)	
	Whorl	40 (19.5)	3.0 (1.0 – 4.0)		22.8 (20.6 – 24.0)	
Ring finger	Arch	14 (6.8)	2.0 (1.0 – 4.0)	0.416	22.0 (21.4 – 23.3)	0.845
	Radial loop	8 (3.9)	2.0 (-0.3 – 2.4)		22.5 (20.1 – 23.9)	
	Ulnar loop	133 (64.4)	3.0 (1.6 – 4.0)		22.0 (20.0 – 24.0)	
	Whorl	50 (24.4)	3.0 (1.0 – 4.0)		22.0 (21.0 – 23.3)	
Fifth finger	Arch	6 (2.9)	2.0 (2.0 – 3.5)	0.019^d	26.0 (22.0 – 27.0)	0.005^c
	Radial loop	2 (1.0)	NA		22.5 (22.0 – 23.0)	
	Ulnar loop	175 (85.4)	2.5 (1.0 – 4.0)		22.0 (20.0 – 23.0)	
	Whorl	22 (10.7)	4.0 (2.8 – 5.0)		23.0 (21.5 – 25.0)	

Kruskal-Wallis test; NA – not applicable (both values were 3.0°)

Table 4b: Post-Hoc pairwise comparison of median values of ANB and FMA between dermatoglyphic patterns of the left hand

Hand	Digit	Paired Dermatoglyphic pattern	Test statistic (ANB)	p-value	Test statistic (FMA)	p-value
Left	Index finger	Arch vs Radial loop	172.0	0.009	181.0	0.016
		Arch vs Whorl	795.0	0.028	934.0	0.252
		Arch vs Ulnar loop	1113.0	0.107	1352.5	0.886
		Radial loop vs Whorl	501.5	0.133	471.5	0.071
		Ulnar loop vs Whorl	2752.0	0.527	2561.0	0.183
		Ulnar loop vs Radial loop	600.5	0.069	476.5	0.004
	Middle finger	Arch vs Radial loop	NA	NA	29.5	0.053
		Arch vs Whorl	NA	NA	611.5	0.257
		Arch vs Ulnar loop	NA	NA	1576.5	0.006
		Radial loop vs Whorl	NA	NA	49.5	0.210
		Ulnar loop vs Whorl	NA	NA	2169.5	0.206
		Ulnar loop vs Radial loop	NA	NA	188.5	0.399
	Fifth finger	Arch vs Radial loop	4.0	0.465	3.0	0.429
		Arch vs Whorl	34.5	0.071	45.5	0.247
		Arch vs Ulnar loop	499.0	0.835	199.5	0.009
		Radial loop vs Whorl	15.0	0.455	19.0	0.752
		Ulnar loop vs Whorl	1148.5	0.002	1284.0	0.010
		Ulnar loop vs Radial loop	144.0	0.664	133.0	0.557

Mann-Whitney U test; NA – Not applicable.

Table 4a: Relationship between participants' dermatoglyphic patterns and quantitative measures of skeletal malocclusion (Contd)

Hand	Digit	Dermatoglyphic pattern	No(%)	ANB (°) Median (IQR)	p-value	FMA (°) Median (IQR) [†]	p-value
Right	Thumb	Arch	23 (11.2)	3.0 (2.0 – 4.0)	0.127	22.0 (21.1 – 24.8)	0.141
		Radial Loop	7 (3.4)	6.0 (2.0 – 9.0)		24.0 (21.3 – 28.3)	
		Ulnar Loop	91 (44.4)	2.8 (2.0 – 4.0)		22.0 (20.6 – 23.0)	
		Whorl	84 (41.0)	3.0 (1.0 – 4.0)		21.8 (20.0 – 24.0)	
	Index finger	Arch	28 (13.6)	2.0 (-0.5 – 3.0)	0.037^a	22.0 (20.0 – 23.9)	0.029^b
		Radial Loop	18 (8.8)	3.5 (2.0 – 5.0)		23.0 (22.0 – 25.0)	
		Ulnar Loop	99 (48.3)	2.5 (1.3 – 4.0)		21.5 (20.0 – 23.0)	
		Whorl	60 (29.3)	3.0 (1.0 – 4.0)		22.0 (20.5 – 24.0)	
	Middle finger	Arch	23 (11.2)	3.0 (2.0 – 5.0)	0.723	22.5 (22.0 – 25.0)	0.026^c
		Radial Loop	1 (0.5)	2.5 (2.0 – 3.0)		21.0 (20.1 – 21.9)	
		Ulnar Loop	155 (75.6)	3.0 (1.0 – 4.0)		22.0 (20.0 – 23.0)	
		Whorl	26 (12.7)	3.0 (1.0 – 4.0)		22.8 (20.6 – 24.0)	
	Ring finger	Arch	8 (3.9)	2.0 (1.0 – 4.0)	0.416	22.0 (21.4 – 23.3)	0.845
		Radial Loop	2 (1.0)	2.0 (-0.3 – 2.4)		22.5 (20.1 – 23.9)	
		Ulnar Loop	138 (67.3)	3.0 (1.6 – 4.0)		22.0 (20.0 – 24.0)	
		Whorl	57 (27.8)	3.0 (1.0 – 4.0)		22.0 (21.0 – 23.3)	
	Fifth finger	Arch	7 (3.4)	2.0 (2.0 – 3.5)	0.019^d	26.0 (22.0 – 27.0)	0.005^e
		Radial Loop	2 (1.0)	NA		22.5 (22.0 – 23.0)	
		Ulnar Loop	180 (87.8)	2.5 (1.0 – 4.0)		22.0 (20.0 – 23.0)	
		Whorl	16 (7.8)	4.0 (2.8 – 5.0)		23.0 (21.5 – 25.0)	

Kruskal-Wallis test; NA¹ – Not applicable (value was -2.0); NA² – Not applicable (FMA value was 20.0); NA³ – Not applicable (ANB value was 2.0).

Table 4b: Post-Hoc pairwise comparison of median values of ANB and FMA between dermatoglyphic patterns of the right hand (Contd)

Hand	Digit	Paired Dermatoglyphic pattern	Test statistic	p-value
Right	Thumb	Arch vs Radial loop	41.5	0.053
		Arch vs Whorl	683.5	0.031
		Arch vs Ulnar loop	579.0	0.001
		Radial loop vs Whorl	261.0	0.621
		Ulnar loop vs Whorl	3518	0.361
		Ulnar loop vs Radial loop	311.5	0.922

Mann-Whitney U test

There were statistically significant differences in the median ANB angle of the participants' fingerprint patterns within the left index and fifth fingers. These differences were noted to exist specifically between the arch and radial loop pattern (p=0.009) and the

arch and whorl pattern (p=0.028) of the left index finger; and the whorl and ulnar loop of the left fifth finger (p=0.009). Using the FMA as a measure of skeletal malocclusion, there were also significant differences in the median values of the left index,

middle, and fifth fingers. The radial and ulnar loops ($p=0.004$) and arch and radial loops of the left index finger; the arch and ulnar loop of the left middle finger ($p=0.006$); and the arch and ulnar loop ($p=0.009$) and ulnar loop and whorl of the left fifth finger ($p=0.010$) all demonstrated statistically significant differences.

The median FMA of the study participants demonstrated significant differences based on their

right thumbs fingerprint patterns only. This difference was particularly common among those with the arch and ulnar loop ($p=0.001$) and the arch and whorl ($p=0.031$) fingerprint patterns. The median ANB values were similar in all the study participants, irrespective of the type of fingerprint pattern they had (see Table 4 above).

Table 5: Association between ABO blood group and skeletal malocclusion

Characteristic	Skeletal Pattern			Total	p-value ¹
	Pattern 1	Pattern 2	Pattern 3		
Blood group A	13 (45.0%)	4 (14.0%)	12 (41.0%)	29 (100.0%)	0.200
Blood group B	11 (45.8%)	2 (8.3%)	11 (45.8%)	24 (100.0%)	
Blood group AB	5 (56.0%)	2 (22.0%)	2 (22.0%)	9 (100.0%)	
Blood group O	85 (59.4%)	26 (18.2%)	32 (22.4%)	143 (100.0%)	
Total	114 (55.6%)	34 (16.6%)	57 (27.8%)	205 (100.0%)	

¹Fisher's exact test

As shown in Table 5 above, most of the individuals with blood group A (45%) had skeletal pattern I malocclusion; skeletal patterns 1 and 2 were equally dominant in individuals with blood group B (46%); Also, most of the individuals with blood group AB

Concerning the Rhesus blood group, most of the individuals who were Rhesus positive (57.0%) had skeletal pattern 1 malocclusion; skeletal pattern 1 and 3 malocclusion were equally dominant in individuals with Rhesus negative (44.4%). There was no

Table 6: Association between Rhesus blood group and skeletal malocclusion

Characteristic	Skeletal Pattern			Total	p-value ¹
	Pattern 1	Pattern 2	Pattern 3		
Rhesus Blood group					0.300
Positive	106 (57.0%)	32 (17.0%)	49 (26.0%)	187 (100.0%)	
Negative	8 (44.4%)	2 (11.1%)	8 (44.4%)	18 (100.0%)	
Total	114 (55.6%)	34 (16.6%)	57 (27.8%)	205 (100.0%)	

¹Fisher's exact test

(56%) as well as blood group O (59%) had skeletal pattern 1 malocclusion. However, there was no statistically significant association between blood group and skeletal malocclusion ($p=0.200$).

statistically significant association between rhesus and skeletal malocclusion ($p=0.300$). Table 6

Table 7: Relationship between participants' blood groups and quantitative measures of skeletal malocclusion

Blood group	No(%)	ANB (°) Median (IQR)	p-value	FMA (°) Median (IQR) [†]	p-value
A	29 (14.1)	2.0 (1.0 – 3.5)		22.0 (20.5 – 23.0)	
B	24 (11.7)	1.5 (-1.0 – 4.0)	0.053	21.3 (20.0 – 23.4)	0.794
AB	9 (4.4)	3.0 (1.5 – 5.0)		22.0 (20.5 – 24.0)	
O	143 (69.8)	3.0 (2.0 – 4.0)		22.0 (20.5 – 24.0)	
Rhesus D pos	187 (91.2)	3.0 (1.5 – 4.0)	0.719§	22.0 (20.5 – 24.0)	0.872§
Rhesus D neg	18 (8.8)	3.0 (1.0 – 4.0)		21.5 (20.4 – 23.3)	

[†]Kruskal-Wallis test; [§]Mann-Whitney U test

Table 8: Correlation of participants' blood groups with categories of skeletal malocclusion

Blood Group	Skeletal pattern 1 <i>N</i> =114 No (%)	Skeletal pattern 2 <i>N</i> =34 No (%)	Skeletal pattern 3 <i>N</i> =57 No (%)	Kendall's tau (τ_b)	p-value
A	13 (11.4)	4 (11.8)	12 (21.1)		
B	11 (9.6)	2 (5.9)	11 (19.3)	-0.143	0.031
AB	5 (4.4)	2 (5.9)	2 (3.5)		
O	85 (74.6)	26 (76.5)	32 (56.1)	0.089	0.225
Rhesus D pos	106 (93.0)	32 (94.1)	49 (86.0)		
Rhesus D neg	8 (7.0)	2 (5.9)	8 (14.0)		

The median (IQR) ANB value in participants with blood group O was 3.0 (2.0 – 4.0) degrees while that of those with blood group B was 1.5 (-1.0 – 4.0). There was, overall, no significant difference between the median values of ANB across the ABO blood groups ($p = 0.053$). The median (IQR) FMA value was similar in the various ABO blood group types as well ($p = 0.794$). The same was true for the ANB and FMA values of the participants across their Rhesus antigen type (Table 7).

When considered in categories, the ABO blood groups tended to occur mostly in the skeletal pattern 1 malocclusion compared to other skeletal

malocclusion patterns. The median (IQR) ANB value in participants with blood group O was 3.0 (2.0 – 4.0) degrees while that of those with blood group B was 1.5 (-1.0 – 4.0). There was overall, no significant difference between the median values of ANB across the blood groups ($p = 0.053$). The median (IQR) FMA value was similar in the various blood group types as well ($p = 0.794$). The same was true for the ANB and FMA values of the participants across their Rhesus antigen type. Skeletal malocclusion related weakly and positively with rhesus antigen positivity, though this was not statistically significant (Table 8).

Table 9: Multivariate analysis of predictors of skeletal malocclusion in study participants

Variable	Adjusted Odds Ratio	95% Confidence Interval	p-value
ABO Blood Group			
A	3.32	0.90 – 12.22	0.071
B	9.62	1.24 – 74.46	0.030
AB	1.13	0.20 – 6.34	0.887
O	1.000		
Left thumb			
Arch	0.44	0.09 – 2.22	0.316
Radial loop	1.57×10^9	0.00 – ∞	0.999
Ulnar loop	0.71	0.23 – 2.15	0.545
Whorl	1.000		
Left index finger			
Arch	0.90	0.26 – 3.10	0.861
Radial loop	1.34	0.24 – 7.51	0.738
Ulnar loop	0.82	0.32 – 2.11	0.685
Whorl	1.000		
Left fifth finger			
Arch	0.92	0.10 – 8.50	0.941
Radial loop	0.00	0.00 – ∞	0.999
Ulnar loop	2.53	1.23 – 5.23	0.012
Whorl	1.000		
Right thumb			
Arch	5.60	0.66 – 47.78	0.115
Radial loop	0.40	0.03 – 5.56	0.498
Ulnar loop	2.75	0.96 – 7.91	0.060
Whorl	1.000		

Further analyses were carried out to explore prediction of skeletal malocclusion in this cohort (Table 9).

Variables that demonstrated a p-value of <0.10 in bivariate analysis were selected and put in a model to predict higher classes of skeletal malocclusion. They included the ABO blood group, left thumb, index and fifth finger and right thumb. After adjusting for confounders, blood group B and an ulnar loop fingerprint pattern on the left fifth finger

independently predicted higher classes of skeletal malocclusion. Participants with blood group B compared to those with group O had about 10 times greater odds of having higher classes of skeletal malocclusion (OR – 9.6 95%CI: 1.2 – 74.5) while possessing an ulnar loop on the left fifth finger compared to possessing a whorl pattern increased the odds of a higher class of malocclusion by 1.5 times (OR – 2.5, 95%CI: 1.2 – 5.2).

Discussion

Genetic and environmental factors play a role in the development of malocclusion that can affect the general health of patients depending on its degree of severity.²⁸ The orofacial structures and the epidermal ridges are derivatives of ectoderm of the embryonic tissue.²⁸ Understanding this biology may play a role in early diagnosis and treatment plan, thereby decreasing the burden of the condition in orofacial region.²⁸

This study assessed the relationship between skeletal malocclusions with fingerprints and blood group as against the background of the role of genetic and environmental factors in the development of malocclusion.

In this study, the most common skeletal pattern was skeletal pattern 1 malocclusion, followed by skeletal pattern 3 and skeletal pattern 2 malocclusion. The most common dermatoglyphic pattern seen among study participants was ulnar loop, followed by whorl, arch and radial loop and is like previous findings in Nigeria by Mohammed et al.,²⁹ in the Kanuri ethnic group. Also, Ujaddughe et al.,³⁰ in their assessment of dermatoglyphic patterns and sex distribution in Esan ethnic group of Edo state, observed that the loop pattern had the highest frequency followed by whorl, arch and double whorl.³⁰ These findings suggest a variable prevalence of fingerprint patterns in Nigerians. Blood group O and Rhesus positive were most predominant among the participants with AB and Rhesus negative being the least prevalent. This was in concordance with other studies done in Nigeria³¹⁻³³, which found Blood group O to be most prevalent and AB to be the least prevalent in the ABO blood group and Rhesus positive to be most prevalent in the Rhesus blood group system.

The dermatoglyphic pattern seen in this study showed that the ulnar loop had the most dominant pattern across all types of skeletal malocclusion, with absence of arch and radial loop patterns on the left fifth fingers in skeletal class 2 and 3 malocclusions. While possessing an ulnar loop on the left fifth finger compared to having a whorl pattern increased the chances of a higher class of malocclusion by 1.5 times. The dermatoglyphic patterns of four fingers out of the ten fingers (left thumb, left index finger, left

fifth finger and Right thumb) had a significant relationship with the different classes of skeletal malocclusion. The ulnar loop pattern of the left fifth finger is associated with skeletal pattern 3, and the left index finger is associated with skeletal pattern 2 malocclusion. The whorl pattern of the left thumb is associated with skeletal pattern 3 malocclusion and whorl pattern of the right thumb is found to be associated with skeletal pattern 2 malocclusion. The left hand had more digits with a significant relationship with skeletal malocclusion. This finding is comparable to that of George et al.¹⁸ and in contrast with the findings of Kaur et al.,³⁴ who found no significant correlation between dermatoglyphics and different skeletal patterns. The contrasting findings may be due to differences in sample size, which was smaller, and the fingerprint method used in the study.³⁴ Reddy et al.,¹⁹ also reported significant association between skeletal malocclusion and specific types of ridge patterns as seen in this study. While different researchers have reported different frequencies of occurrence of specific dermatoglyphic patterns with different classes of skeletal malocclusion. It is understandable that their divergent findings may be due to differences in population studied among other factors because the cohort in this present study consisted of those with some degree of malocclusion and a relatively uniform population.

The bivariate analysis of blood groups with skeletal malocclusion did not establish any relationship in this study. However, the multivariate analysis shows that blood group B when compared with other blood groups was found to be directly related to higher classes of skeletal malocclusion. This aligns with the findings by Gheisari et al.³⁵ which revealed that blood groups B and AB have an increased incidence of association with maxillofacial deformities, whereas, blood groups O and A have a lower incidence of association in Iranian populations. A contrary opinion was, however, held by Shokor et al.,³⁶ who reported no genetic influence of ABO blood group in relation to variations in craniofacial morphology. The reason for this may be due to the difference in the study methodologies.

The dermatoglyphic patterns of individuals remain constant throughout life, from birth till death and have

been found to have a relationship with malocclusion, although, it requires more research to be done. This easy, accessible, inexpensive, useful, reliable, and non-invasive method may be a useful tool if employed in clinical settings to aid early assessment of risk for malocclusion, along with diagnostic impression of sagittal skeletal relationships of maxilla and mandible in malocclusion. This might aid early preventive and interceptive orthodontic management of malocclusion if established, especially in developing countries with enormous populations and limited health budgets. In addition, the practice of routine recording of the fingerprints of patients in the orthodontic clinic along with other orthodontic records will go a long way in assisting forensic investigations and anthropology studies. The sample size was a limitation, the participants were all hospital patients. No participants with normal occlusion were recruited in the study and this led to some assumptions in the regression model.

Conclusion

This study was able to show that there is a relationship between dermatoglyphic pattern with skeletal malocclusions. The possession of ulnar loop on the left fifth finger compared to the possession of whorl pattern increased the odds of a higher class of malocclusion by 1.5 times. There is a marked absence of the arch and radial loop patterns on the left fifth fingers in skeletal class 2 and 3 malocclusions. The ulnar loop patterns of left fifth finger and left index finger were associated with skeletal pattern 3 and

skeletal pattern 2 malocclusion respectively. The whorl pattern of the left thumb was associated with skeletal pattern 3 malocclusion while the whorl pattern of the right thumb was found to be associated with skeletal pattern 2 malocclusion. There was no significant association between blood group and skeletal malocclusion. However, possession of blood group B, when compared with other blood groups, was found to be directly related to higher classes of skeletal malocclusion.

The dermatoglyphic patterns of some digits show predictive values, this relationship could be further studied in more races and in larger populations before being employed as a diagnostic aid for identifying malocclusion problems at an early age for timely orthodontic intervention.

Recommendation:

The dermatoglyphic patterns show the promise of being a predictive tool in skeletal malocclusion, thus, more research in different races and in larger sample populations is needed to further determine the extent of its use. The inclusion of participants with normal occlusion in future studies will aid comparison as well as test the specificity and sensitivity of dermatoglyphic patterns as a tool.

Authors' Contribution: All the authors contributed to the manuscript

Financial Support - Funded by the authors

Conflict of Interest - None to declare

References

- 1 Kaur J, Khokhar V and Pathak A. Dermatoglyphics-as a diagnostic tool in detection of malocclusion. *Int J Rec Sci Res* 2016;7:12400–12404.
- 2 Daskalogiannakis J: Glossary of Orthodontic Terms. Publisher: Quintessence, 2000, 1st Ed.
- 3 Houston W, Stephens C, Tulley W. A textbook of orthodontics. Bristol: Wright Pub Co, 1992.
- 4 Sharma R, Preethi PN, Nagarathna C, Navin HK. "Association of ABO blood groups with malocclusion in the population of Jaipur, India: A prospective study." *Int J Sci Study*. 2015; 2: 45-51.
- 5 Rasool G, Shah SS, Adil M. Relationship of the skeletal malocclusion with dental malocclusion in the sagittal plane. *JKCD* 2016;6: 16-19.
- 6 Nishio C, Huynh N. Skeletal Malocclusion and Genetic Expression : An Evidence- Based Review. *J Dent Sleep Med* 2016;3:57–63.
- 7 Adnan H, Koshak H. Diagnosis , Complications , And Treatment of Dentoskeletal Malocclusion ; Systematic Review. *Int J Healthcare Sci* 2017;5:95–101.

- 8 Divyashree, Suhas AS, SharmadaBK, Tayeepriyanka. Dermatoglyphic Patterns and Their Correlation with Skeletal Malocclusions. *J Dent Med Sci*. 2016;15:101–104.
- 9 Achalli S, Patla M, Nayak U, Soans C R. Dermatoglyphics and orthodontics. *Int J Orthod Rehabil* 2016;7:144-147.
- 10 Gheisari R, Ghoreishian M, Movahedian B, Roozbehi A. The association between blood groups and maxillofacial deformities. *Indian J Plast Surg* 2008;41:138-40
- 11 Fathin FBAS, Wan SWR, Mohammad KA. Craniofacial morphology with genetic influence of ABO blood group in Malaysian orthodontic patients. *Int J Pharm Bio Sci* 2015; 6: 412-418.
- 12 Gupta SP. Association of ABO Blood Groups with Occlusal Pattern among Orthodontic Patients of Kathmandu District. *EC Dent Sci*(2018): 17;2119-2129.
- 13 Ruhl CM, Bellian KT, Van Meter BH, et al. Diagnosis, complications, and treatment of dentoskeletal malocclusion. *Am J Emerg Med*. 1994; 12:98-104.
- 14 Carvalho AC, Paiva SM, Viegas CM, Scarpelli AC, Ferreira FM, Pordeus IA. Impact of malocclusion on oral health-related quality of life among Brazilian preschool children: a population-based study. *Braz Dent J* 2013; 24: 655-661
- 15 Zere E, Chaudhari PK, Sharan J, Dhingra K, Tiwari N. Developing class iii malocclusions: challenges and solutions. *Clin, Cosmetic Investig Dent* 2018;10: 99–116
- 16 Eslami N, Jahanbin A, Ezzati A, Banihashemi E and Kianifar H: Can dermatoglyphics be used as a marker for predicting future malocclusions? *Electr Physician*. 2016; 8:1927–1932.
- 17 Jindal G. A comparative evaluation of dermatoglyphics in different classes of malocclusion. *Saudi Dent J* [Internet]. King Saud University; 2015;27:88–92. Available from: <http://dx.doi.org/10.1016/j.sdentj.2014.11.012>.
- 18 George SM, Philip B, Madathody D, Mathew M, Paul J, Dlima JP. An Assessment of Correlation between Dermatoglyphic Patterns and Sagittal Skeletal Discrepancies. *J Clin. Diagn. Res*. 2017; 11:35 - 40.
- 19 Reddy BRM, Sankar SG, Roy E.T, Govulla S. A Comparative Study of Dermatoglyphics in Individuals with Normal Occlusions and Malocclusions. *J Clin Diag Res*: 2013; 7:3060 - 3065.
- 20 Al-Khatieeb MM, Al-Joubori SK and Taha SS. Association of ABO Blood Group and Rhesus Factor with Dental Malocclusion in a Population of Baghdad, Iraq. *Int J Med Res Health Sci* 2018, 7: 165-169
- 21 Bhasin MT, Bhasin P, Singh A, Bhatia N, Shewale AH, Gambhir N, et al. Dermatoglyphics and malocclusion-A forensic link. *Br Biotec J*, 2016;13:1–12.
- 22 Gravely JF and Johnson DB. Angle's Classification of Malocclusion: an Assessment of Reliability, *Br J Orthod* 2016; 1:3, 79-86,)
- 23 Lakshmi V. Dermatoglyphics and orthodontics - A review. *Ann Essences Dent* 2013; 4:30-33.
- 24 Isiekwe, MC. and Sowemimo GC. Cephalometric findings in a normal Nigerian - population sample and adult Nigerians with unrepaired clefts. *Cleft Palate J* 1984. 21:323–328.
- 25 Steiner CC. Cephalometrics for you and me. *Am J Orthod*, 1953;39: 729-55.
- 26 Proffit, W. R., Fields, H. W., and Sarver, D. M. (2007). *Contemporary orthodontics* (4th ed.). St. Louis, Mo.: Mosby Elsevier.
- 27 Harsha L, Mds GJ. Correlation of Lip Print , Fingerprint and Blood Groups in a Tamil Nadu Based Population . *J. Pharm. Sci. and Res* 2015;7:795–799.
- 28 Prakash P, Sirjana D, Vivek B T, Amrita S, Prabesh S Dermatoglyphic Pattern and Types of Malocclusion among Individuals visiting A Medical Institution of Nepal. *Int J Orthod Rehabil* 2020;20:46-50.
- 29 Mohammed, BS. Garba, SH and Adeyemi LB .Digital Dermatoglyphics Patterns of the Kanuri Ethnic Group of North Eastern Nigeria. *Int J Innovation and Applied Studies* .2014; 9: 985-988.
- 30 Ujaddughe MO, Abue AD, Izunya MA, Ezeuko

- VC, Eze IG. and Baxter-Grillo D. assessment of dermatoglyphic patterns and sex distribution in Esan ethnic group of Edo state, Nigeria. IJBAIR, 2015, 4: 9 - 14.
- 31 Enosolease ME, Bazuaye GN. Distribution of ABO and Rh-D blood groups in the Benin area of Niger-Delta: Implication for regional blood transfusion. Asian J Transfus Sci. 2008 ; 2: 3–5.
- 32 Odokuma EI, Okolo AC, Aloamaka PC. Distribution of ABO and rhesus blood groups in Abraka, Delta State, Nigeria. Niger J Physiol Sci. 2007; 22:89-91.
- 33 Kooffreh ME, Ikpeme EV, Ekerette EE, Eyo NO. Evaluating the Prevalence of Five Genetic Traits of Simple Inheritance in Association with the Distribution Pattern of ABO and Rhesus Phenotypes among Families in Calabar, Nigeria. J Med Sci. 2015; 15: 185-191.
- 34 Kaur H, Tikku T, Khanna R, Maurya RP, Verma S, Srivastava K, et al. Assessment of correlation between dermatoglyphics of individuals with different skeletal growth. Int J Orthod Rehabil 2020;11:69-75.
- 35 Gheisari R, Ghoreishian M, Movahedian B, and Roozbehi A. The association between blood groups and maxillofacial deformities. Indian J Plast Surg. 2008 41: 138–140.
- 36 Shokor FFBA, Rahman WSWA and Alam MK craniofacial morphology with genetic influence of ABO blood group in Malaysian Orthodontic patients. Int J Pharm Bio Sci 2015 ; 6: 412-41

Correlations between Cephalometric Measurements of Hard Tissues and Photogrammetric Features of Facial Soft Tissues.

Djibril Cisse,* Alpha Badiane, Malick Guèye, Irado Rakoto Solofo, Joseph Samba Diouf, Khady Diop-Bâ, Papa Ibrahima Ngom .

Abstract

Background: Facial soft tissues cover the teeth and the facial skeleton; maintain close anatomical relations with them and may nevertheless vary in their morphology and position. The objective of this study was to determine the correlations between soft facial tissue measurements and craniofacial hard tissue measurements..

Methods: A cross-sectional descriptive study was performed using standardised tele-radiographic and photographic images of young children who had come for orthodontic care. On each selected subject, cephalometric and photogrammetric measurements were made. The data collected was analysed using the IBM SPSS 20.0 statistical software. The correlation between hard tissue cephalometric measurements and photogrammetric soft tissue measurements was investigated by a Pearson correlation. The significance is fixed at $p=0.05$.

Results: ANB was significantly and positively correlated to Sn-N'-Sm. I / A-Pog was significantly and positively correlated with Ls-E and Li-E. Significant and positive correlations were found between the position of the pogonion and Sn-H. It is the same between the position of point A and Trg-Sn distance; and between S-Ar and the variables N'-Sn and N'-Me'. Ar-Go was also significantly and positively correlated to Sn-Me', Sn-Sts and N'-Me'.

Conclusion: The many significant and positive correlations found between cephalometric and photogrammetric variables show that these could advantageously be used to evaluate skeletal and dentoalveolar variables. This would limit the significant risks of ionizing radiation during orthodontic treatment.

Keywords: Photogrammetry, cephalometry, hard tissue, soft tissue

Authors' affiliations

Orthodontic Department/Service, Department of Odontology,
Faculty of Medicine, Pharmacy and Odontology,
Cheikh Anta DIOP University, Dakar, Senegal

Correspondence:

Djibril Cisse
BP 25247 Dakar, Fann, Senegal.
Email: djibrilcisse3@yahoo.fr

Introduction

The photogrammetry of the soft tissues of the face is a measurement method that objectively quantifies the characteristics of the face through the photographic tool. Researchers have used it to evaluate the characteristics of the face^{1,2} Unlike cephalometry, it is inexpensive and does not expose the patient to ionizing radiation. In addition, it is reliable and sufficiently reproducible because it is easy to achieve in a conventional context, without the need for special equipment.¹ Facial soft tissues cover the teeth and the facial skeleton, maintain close

anatomical relationships with them and can vary considerably in their morphology and position. These variations may be related to the existence of inter-ethnic, racial and geographical variability in the facial skin tissues² but also to the existence of a positional and morphological variability of the underlying skeletal and dentoalveolar structures.³ The formulation of a treatment plan based solely on cephalometric measurements of skeletal and dentoalveolar structures is therefore insufficient and often leads to aesthetic problems.^{4,7} Therefore, it is necessary to have these cephalometric measurements coupled with photogrammetric measurements in order to better identify the skeletal and dentoalveolar abnormalities on the one hand and the cutaneous ones on the other, and their possible associations. However, little importance has been devoted to quantifying the relationships between the position or dimensions of the components of soft facial tissues and different skeletal and dentoalveolar parameters.

Most research in this area focuses on the study of cephalometric and photogrammetric projection and identification errors⁸ and the response of soft tissues to the movements of orthopedic, orthodontic or orthognathic therapy.⁹⁻¹²

Separate cephalometric and photogrammetric measurements have been performed in a Senegalese population,^{13,14} but no study of associations between these two parameters has been performed in this population. The objective of this study was to determine the correlations between the photogrammetric measurements of the face and different profile teleradiographic measurements in Senegalese subjects.

Materials and methods

It is a transversal analytical study carried out at the orthodontic clinic of the Department of Odontology of Cheikh Anta Diop University in Dakar. The study is based on a sample of a group of Senegalese children aged between 6 and 12 who came for consultation to the Dentofacial Orthopedics Clinic of the Department of Odontology of the Faculty of Medicine, Pharmacy and Dentistry of the Faculty of Medicine, Pharmacy and Dentistry of Cheikh Anta Diop University in Dakar. Subjects who received orthodontic or prosthetic treatment, plastic or orthognathic surgery were not included in the study. It is the same for those who had orofacial soft tissue pathologies (swelling, ulcerations ...) or severe craniofacial abnormalities. The pictures on which one could see a contracture of the muscles of the chin tassel (pleated or flattened chin) were also excluded from the study.

Photogrammetric recordings and measurements:

On each subject selected according to the above criteria, a standardised profile photographic snapshot is made according to the method described by Ferrario et al.¹⁵ All photos were taken with the same digital camera (Samsung type, with a resolution of 14.2 megapixels, optical zoom $\times 3$). The images obtained were digitised on a computer using a photo software (photoshop type elements 7.0) with a resolution of 300 DPI (dot per inch or PPP point par pouce). These scanned images were printed on a

white sheet from a single printer (HP Deskjet 3050). The magnification of the images obtained was calculated from that of the image of a metric scale taken as a reference. From that moment, the actual linear measurements could be obtained. The points and lines used for this approach are defined in Table I. The photogrammetric variables are defined in Table II and illustrated in Figures 1 and 2.

Cephalometric recording: Cephalometric measurements were then made from a teleradiographic profile of each selected subject. Informed consent was obtained from the parents before the snapshot was taken. Plotting all the teleradiographies was done manually by a single examiner (JD) on an acetate paper sheet. The points and lines chosen are defined in Table 1. The measured cephalometric variables are defined in Table II and anatomically illustrated in Figures 3 and 4.

Statistical analyses: The data collected was analysed using the IBM SPSS 20.0 statistics software for windows. The power of the association between photogrammetric measurements and cephalometric variables was sought by the Pearson correlation coefficient. The materiality threshold is set at $p = 0.05$.

Method error: The reliability of photogrammetric and cephalometric measurements was assessed by measuring again, at random, the variables in a group of 30 children (representing 30% of study subjects), one month after the first assessment. The Cronbach test was performed to determine the error of the method. An intra-class coefficient between 0.789 and 0.971 was found. This result shows that there is no significant difference between the first and second measurements.

Results: One hundred subjects comprising 49 boys and 51 girls aged between 6 and 12 years with an average age of 9.5 ± 1.08 years were included in this study.

Correlations between Sagittal Photogrammetric Variables and Sagittal Teleradiographic Parameters

The position of point A was significantly and positively correlated with Trg-Sn ($r = 0.40$ and $p =$

0.04) and Sn-N'-Sm ($r = 0.52$ and $p = 0.008$). I/A-Pog was also significantly and positively correlated with Ls-E ($r = 0.51$ and $p = 0.01$), Li-E ($r = 0.48$ and $p = 0.01$), Ls-S ($r = 0.47$, $p = 0.01$) and Li / Sn-Sm ($r = 0.46$, $p = 0.02$). The convexity was significantly and positively correlated with Sn-N'-Sm ($r = 0.41$ and $p = 0.04$). It is the same between the position of the pogonion and the Sn-H photogrammetric variable ($r = 0.42$, $p = 0.03$). The ANB angle was also significantly and positively correlated with the Sn-N'-Sm angle ($r = 0.52$ and $p = 0.009$) (Table 3).

Correlations between vertical photogrammetric variables and vertical teleradiographic parameters

The N-Me height was significantly and positively correlated with Sn-Me' and N'-Sn distances with r equal to 0.46 and 0.56 respectively; and p respectively equal to 0.02 and 0.004. This N-Me height was also significantly and positively correlated with Sti-Sm and N'-Me' with r respectively equal to 0.66 and 0.67 and p values < 0.001 . N-ENA was significantly and positively correlated to N'-Sn, Sti-Sm and N'-Me' with r respectively equal to 0.51; 0.55 and 0.56; and p respectively equal to 0.01; 0.001

and 0.004. ENA-Me was significantly and positively correlated with Sn-Me' ($r = 0.48$ and $p = 0.01$), N'-Sn ($r = 0.51$ and $p = 0.01$), Sn-Sts ($r = 0.45$ and $p = 0.02$), Sti-Sm ($r = 0.55$ and $p = 0.005$) and N'-Me' ($r = 0.66$ and $p < 0.001$). S-Go was significantly and positively correlated to the distances N'-Sn, Sn-Sts, Sti-Sm and N'-Me' with r respectively equal to 0.59; 0.41; 0.65 and 0.61 and p respectively equal to 0.002; 0.04; 0.001 and 0.001. The S-Ar height was significantly and positively correlated with N'-Sn distances ($r = 0.63$ and $p = 0.001$), Sti-Sm ($r = 0.51$ and $p = 0.009$) and N'-Me' ($r = 0.55$ and $p = 0.005$). The Ar-Go distance was significantly and positively correlated with Sn-Me', N'-Sn, Sn-Sts, Sti-Sm, Sm-Me' and N'-Me' with r respectively equal to 0.41; 0.41; 0.48; 0.65; 0.42 and 0.55 and p respectively equal to 0.04; 0.04; 0.01; 0.001; 0.03 and 0.005. The Y axis was significantly and positively correlated with the Sm-Me' / Li-Sm angle ($r = 0.40$ and $p = 0.04$) and negatively correlated with angle Z ($r = -0.54$ and $p = 0.006$). The gonadal angle was significantly and positively correlated with Sm-Me' / Li-Sm ($r = 0.43$ and $p = 0.03$) (Table 4).



Figure 1: Sagittal photogrammetric variables: 1 = Trg-Sn; 2 = Sn-H; 3 = Ls-E; 4 = Ls-S; 5 = Ls / Sn-Sm; 6 = Li-S; 7 = Li-E; 8 = Li / Sn-Sm; 9 = Pog' / Sn-Sm; 10 = Sn-N'-Sm.



Figure 2: vertical photogrammetric variables: 1 = Sn-Me'; 2 = Sn-Sts; 3 = Sti-Sm; 4 = Sm-Me'; 5 = N'-Sn; 6 = N'-Me'; 7 = Z-angle; 8 = Prn-Sn-Ls; 9 = Sn-Ls / Li-Sm; 10 = Sm-Me' / Li-Sm.




Figure 3: sagittal cephalometric variables: 1 = I / F; 2 = ANB; 3 = position of point A; 4 = convexity; 5 = I / A-Pog; 6 = sagittal position of the pogonion point; 7 = i / M.



Figure 4: Vertical cephalometric variables: 1 = N-ENA; 2 = ENA-Me; 3 = N-Me; 4 = Y axis; 5 = S-Go; 6 = S-Ar; 7 = Ar-Go; 8 = angular / bony angle; 9 = FMA.

Table 1: Definition of points and lines used

Skin points and lines	Definitions
Nasion (N')	the point of the midline at the root of the nose
Pronasal (Prn)	the most salient point of the tip of the nose
Sub nasal (Sn)	the point where the upper lip joins the columella
Upper labial (Ls)	the point that indicates the mucocutaneous limit of the upper lip
Upper Stomion (Sts)	the lowest point of the upper lip
Lower stomion (Sti)	the highest point of the lower lip
Lower labial (Li)	the point that indicates the mucocutaneous border of the lower lip
Supramental (Sm)	the deepest point of the sub-labial lower concavity
Pogonion (Pog ')	the most anterior point of the chin
Chin (Me ')	the lowest point of the lower edge of the chin
Tragus (Trg)	the most posterior point of the atrial tragus
Suborbital (Or ')	perceptible point on the finger of the outer orbital rim
Line H	Line passing through the skin chin point and tangent to the upper lip
Line Sn-Sm	Aesthetic line of Canut, right joining Sn and Sm points
(Trg-Or ')	Frankfurt horizontal cutaneous plane, joining tragus and under orbital points
Line E	Ricketts E aesthetic line tangent to the tip of the nose and the tip of the chin
Line S	Line S is the line joining the point Pog and the middle between Prn and Sn
Line Z	line tangent to the chin and the most prominent lip
Skeletal and dentoalveolar points	
Nasion (N)	anteroposterior point of the image of the naso-frontal suture
Sella Turcica (S)	Sella Turcica (S)
Articulare (Ar)	point of the image of the posterior border of the ramus with the exocranial face of the occipital clivus
Porion (Po)	highest point of the external auditory canal located on the vertical passing through its middle
Anterior nasal spine (ANS)	tip of the anterior nasal spine
A	most sloping point of the concavity of the maxillary image
Orbital (Or)	lowest point of the orbit image
Upper Incisor (Ui)	free edge of the most anterior superior incisor
B	most sloping point of the concavity of the image of the symphysis
Chin (Ch)	lowest point of the image of the mandibular symphysis
Gonion (Go)	point equidistant from the most posterior point of the horizontal branch of the mandible, and from the lowest point of the rising branch
Pogonion (Pog)	most salient point of the image of the symphysis
Gnathion (Gn)	equidistant point between the point Pog and Me
Basion (Ba)	lowest point of the occipital basi

Table 2: Definition of chosen variables

Cephalometric variables	Definitions
I / A-Pog	distance from the incisal edge of the medial incisor superior to line A-Pog.
Sagittal position of the pogonion	distance between the point pogonion and the perpendicular to the Frankfurt plane passing through point Na.
Position of point A	distance between A and the perpendicular to PHF passing through point N
Convexity	distance from A to Na-Pog facial plane
I / F	angle that makes the axis of the upper incisor with the horizontal plane of Frankfurt
i / M	angle made by the axis of the lower incisor with the mandibular plane
ANB	maxillomandibular sagittal shift
N-Me	Total anterior facial height
N-ENA	upper anterior facial height
ENA-Me	lower anterior facial height
S-GB	total posterior head height
S-Ar	superior posterior head height
Ar-Go	lower posterior facial height
FMA	angle between the Frankfurt plane and the Downs mandibular plane
1- Y axis	angle formed by the line (SGn) and the Frankfurt plane
2- Gonic / Angular / Bony angle	angle formed by the mandibular plane (GoMe) and the tangent to the posterior edge of the mandible (ArGo).
1- Photogrammetric variables	
Trg- Sn	1- depth of the face
Ls-E	distance from the upper lip Ls to line E
Li-E	distance from lower lip Li to line E
Sn-H	distance from sub nasal point Sn to line H
Ls-S	distance from Ls to line S
Li-S	Li's distance to S line
Ls/ Sn- Sm	distance from Ls to Canut line (Sn-Sm)
Li/ Sn-Sm	distance from Li to Canut line (Sn-Sm)
Pog'/Sn- Sm	distance from Pog to Canut line (Sn-Sm)
Sn- N'- Sm	cutaneous ANB angle
Sn-Me'	lower third of the face
N'-Sn	nose length
Sn-Sts	length of the upper lip
Sti-Sm	length of the lower lip
Sm-Me'	chin height
N'-Me'	height of the lower floor
Prn-Sn-Ls	naso labial angle
Sn-Ls/Li-Sm	Inter labial angle
Sm-Me'/Li-Sm	Mento-labial angle
Z angle	formed by the intersection of the plane Frankfurt horizontal cutaneous (Trg-Or ') and line Z

Table 3: Correlations between sagittal photogrammetric variables and sagittal telerradiographic parameters

Settings Parameters (mm,°)	I/A-Pog r (p)	Position Pog r (p)	Position A r (p)	Convexit� r (p)	I/F r (p)	i/M r (p)	ANB r (p)
Trg-Sn	0.16 (0.43)	0.33 (0.11)	0.40* (0.04)	0.25 (0.22)	0.04 (0.83)	0.05 (0.81)	0.19 (0.35)
Ls-E	0.51* (0.01)	-0.22 (0.28)	0.09 (0.66)	0.36 (0.07)	-0.13 (0.52)	-0.02 (0.89)	0.25 (0.23)
Li-E	0.48* (0.01)	-0.36 (0.08)	-0.24 (0.25)	0.11 (0.60)	0.03 (0.88)	-0.26 (0.20)	-0.03 (0.86)
Sn-H	-0.28 (0.17)	0.42* (0.03)	0.24 (0.25)	-0.14 (0.50)	0.21 (0.32)	0.14 (0.51)	-0.03 (0.85)
Ls-S	0.47* (0.01)	-0.08 (0.68)	0.18 (0.39)	0.31 (0.13)	-0.07 (0.72)	0.01 (0.94)	0.26 (0.20)
Li-S	0.34 (0.10)	-0.39 (0.05)	-0.36 (0.08)	-0.02 (0.91)	-0.13 (0.53)	-0.16 (0.44)	-0.12 (0.55)
Ls/ Sn- Sm	0.14 (0.50)	0.19 (0.36)	0.39 (0.05)	0.25 (0.23)	-0.10 (0.61)	0.18 (0.38)	0.35 (0.09)
Li/ Sn-Sm	0.46* (0.02)	-0.1 (0.64)	-0.20 (0.33)	-0.04 (0.82)	0.11 (0.6)	-0.18 (0.39)	-0.10 (0.63)
Pog'/Sn- Sm	-0.01 (0.93)	0.30 (0.14)	0.09 (0.66)	0.11 (0.58)	0.19 (0.36)	0.08 (0.68)	0.01 (0.96)
Sn- N'- Sm	-0.08 (0.70)	0.28 (0.18)	0.52** (0.008)	0.41* (0.04)	-0.08 (0.70)	-0.03 (0.87)	0.52** (0.009)

*= The correlation is significant at the 0.05 level (bilateral). ** = The correlation is significant at the 0.01 level (bilateral).

Table 4: Correlation between vertical photogrammetric variables and vertical telerradiographic Parameters

Parameters (mm,°)	N-Me r (p)	N-ENA r (p)	ENA-Me r (p)	S-Go r (p)	S-Ar r (p)	Ar-Go r (p)	FMA r (p)	Y Axis r (p)	Angular / Bony angle r (p)
Sn-Me'	0.46* (0.02)	0.39 (0.05)	0.48* (0.01)	0.38 (0.06)	0.25 (0.23)	0.41* (0.04)	-0.18 (0.4)	-0.11 (0.60)	0.17 (0.40)
N'-Sn	0.56** (0.004)	0.49* (0.01)	0.51* (0.01)	0.59** (0.002)	0.63** (0.001)	0.41* (0.04)	-0.19 (0.37)	0.02 (0.91)	-0.01 (0.92)
Sn-Sts	0.33 (0.11)	0.17 (0.41)	0.45* (0.02)	0.41* (0.04)	0.08 (0.69)	0.48* (0.01)	-0.10 (0.62)	0.07 (0.73)	-0.07 (0.74)
Sti-Sm	0.66** (<0.001)	0.62** (0.001)	0.55** (0.005)	0.65** (0.001)	0.51** (0.009)	0.65** (0.001)	-0.23 (0.26)	0.19 (0.35)	-0.20 (0.34)
Sm-Me'	0.38 (0.06)	0.30 (0.15)	0.40 (0.05)	0.33 (0.10)	0.25 (0.23)	0.42* (0.03)	0.01 (0.96)	0.18 (0.39)	0.27 (0.19)
N'-Me'	0.67** (<0.001)	0.56** (0.004)	0.66** (<0.001)	0.61** (0.001)	0.55** (0.005)	0.55** (0.005)	-0.18 (0.37)	0.02 (0.91)	0.08 (0.70)
Prn-Sn-Ls	0.08 (0.68)	-0.08 (0.68)	0.27 (0.19)	0.02 (0.89)	0.07 (0.74)	-0.03 (0.85)	0.39 (0.05)	0.29 (0.16)	-0.04 (0.83)
Sn-Ls/Li-Sm	-0.07 (0.74)	-0.12 (0.55)	0.01 (0.96)	-0.06 (0.75)	-0.09 (0.67)	-0.001 (0.99)	0.15 (0.46)	0.09 (0.67)	0.12 (0.55)
Sm-Me'/Li-Sm	-0.08 (0.71)	-0.11 (0.58)	-0.04 (0.85)	-0.07 (0.74)	-0.18 (0.37)	0.17 (0.42)	0.28 (0.17)	0.40* (0.04)	0.43* (0.03)
Z Angle	0.17 (0.42)	0.25 (0.23)	0.10 (0.63)	-0.04 (0.84)	-0.04 (0.83)	-0.09 (0.66)	-0.37 (0.07)	-0.54** (0.006)	-0.33 (0.11)

* = The correlation is significant at the 0.05 level (bilateral). ** = The correlation is significant at the 0.01 level (bilateral).

Discussion

The position of point A was significantly and positively correlated with Trg-Sn ($r = 0.40$ and $p = 0.04$). In other words, the deeper the face was, the closer point A was to the line perpendicular to the Frankfurt plane passing through the Nasion point. This correlation could be related to the fact that the advanced point A located on the maxillary alveolar rampart can cause that of the Sn point of the upper lip which rests on this alveolar rampart. When Sn moves forward, the Trg-Sn distance increases. Similarly, when A moves forward, its distance to the line perpendicular to the Frankfurt plane passing through the Nasion point decreases but the variable of the position of A increases since it is negatively noted.

The significant and positive correlations between the position of point A and Sn-N'-Sm on the one hand and between the convexity and Sn-N'-Sm on the other hand can be related to the simultaneous displacement of points A and Sn with the displacement of the maxillary alveolar rim and the upper lip which is overlying it. The further A point moves forward, the more its sagittal position increases as well as the convexity. Also, the more points Sn moves forward, the more the Sn-N'-Sm angle increases; hence the significant and positive correlations between these parameters. The value of the correlation coefficient between the position of point A and the Sn-N'-Sm angle of 0.52 shows a broad association between these two parameters; where the correlation coefficient between the convexity and Sn-N'-Sm of 0.41 (<0.5) shows a mean association according to Cohen [16].

Significant and positive correlations between I / A-Pog and Ls-E and Ls-S variables could also be explained by the relationship between the upper incisor and the upper lip. Several studies have shown that a retroversion of the upper incisor can lead to superior prostration [11,17]. Moreover, the orthodontic treatment of retraction of the upper incisors is often associated with a retraction of the upper lip which concomitantly corrects the prostration. The fact that the upper incisor also rests on the upper part of the lower lip and flush with its free edge explains the significant and positive correlation between the variable I / A-Pog which indicates the degree of superior incisal protrusion and

the Li-E distances ($r = 0.48$ and $p = 0.01$) and Li / Sn-Sm ($r = 0.46$, $p = 0.02$).

The position of the pogonion was significantly and positively correlated with the Sn-H distance ($r = 0.42$, $p = 0.03$). In fact, the distal displacement of the pogonion leads to that of the cutaneous chin. This induces a rotation of the H line around the most prominent point of the upper lip, which means a recoil of the segment located under the center of rotation and an advance of the segment located above the center of rotation. An increase in the Sn-H distance will follow (which means a decrease of this negatively noted variable); hence the significant and positive correlation between the position of the pogonion and the Sn-H variable.

The significant and positive correlation between the ANB and Sn-N'-Sm angles show that variations in maxillomandibular skeletal shift follow that of the maxillomandibular cutoff. Indeed, the Sn-N'-Sm angular photogrammetric variable represents the cutaneous ANB angle. The value $r > 0.5$ shows a wide association between maxillomandibular skeletal shift and maxillomandibular cutoff [16]. This strong association is justified by the fact that the different points which constitute these two angles are situated opposite one another. In fact, cephalometric point A is located opposite cutaneous point Sn. The same is true of the Nasion point (located at the level of the naso-frontal suture) and the Nasion point of the skin at the level of the nasal trellis. Point B (the most sloping point of the mandibular alveolar rim) is also opposite point Sm situated at the bottom of the labial sulcus. Similar results have been reported by Gomes & al. in a study involving a sample of 123 subjects, including 65 girls and 58 boys aged 7 to 12, with whom telerradiographic images and photographic images in profile were taken. The results of this study showed a significant and positive correlation between skeletal ANB and cutaneous ANB ($r = 0.82$, $p < 0.001$) [18]. Adwani & al. also found similar results for a sample of 150 subjects, including 64 boys, and 86 girls aged 13-28, where r and p were 0.689 and 0.001, respectively [19].

Correlation between vertical photogrammetric variables and vertical telerradiographic parameters

The cephalometric variables N-Me, N-ENA, ENA-Me, S-Go, S-Ar and Ar-Go were all significantly and positively correlated with photogrammetric variables

N'-Sn, Sti-Sm and N'-Me'. In addition, the three variables N-Me, ENA-Me and Ar-Go were significantly and positively correlated with Sn-Me '. It is the same between the ENA-Me, S-Go and Ar-Go variables and the Sn-Sts photogrammetric parameter. Specifically, the Ar-Go variable was significantly and positively correlated with the Sm-Me photogrammetric variable. These significant and positive correlations between vertical linear teleradiographic variables and vertical linear photogrammetric variables show the close relationships between skeletal structures and their vertical cutaneous structures. Thus, vertical linear photogrammetric variables could be used to evaluate the linear variables of the skeletal structures in the vertical direction. The use of correlation coefficients makes it possible to obtain the value of cephalometric measurements from photogrammetric measurements. This contributes to a considerable reduction of the risks of irradiation of the subjects when it comes to evaluating the skeletal structures in the vertical direction.

Axis Y was significantly and positively correlated with Sm-Me ' / Li-Sm ($r = 0.40$ and $p = 0.04$) and negatively correlated with angle Z ($r = -0.54$ and $p = 0.006$). The gonadal angle was significantly and positively correlated with Sm-Me ' / Li-Sm ($r = 0.43$ and $p = 0.03$).

The significant and positive correlations between the Y axis (Frankfurt / S-Gn plane) and Sm-Me ' / Li-Sm on the one hand and between the gonionic angle and Sm-Me ' / Li-Sm on the other hand could be related to the concomitant displacement of the Gnathion points, Skeletal Chin and Cutaneous Chin all located in the chin area. Indeed, a recoil of the chin leads to that of

the Gnathion points, skeleton chin and cutaneous chin, those which cause an increase in the angle of the Y axis and the Sm-Me ' / Li-Sm angle as well as an increase in the angle of the glands; hence these significant and positive correlations. The significant and negative correlation between the angle of the Y axis and the Z angle (between the Frankfurt plane and the straight line connecting the Chin point to the most prominent lip) can also be related to the displacement in the same direction of Gnathion and Chin points. A shift of the skin point towards the back causes a reduction of angle Z. On the other hand, a shift of the gnathion point towards the back causes an increase in the angle Y axis, which justifies the significant and negative correlation between the angle of the Y axis and angle Z.

Conclusion

These numerous correlations prove the close relationship between hard facial tissues and their overlapping skin tissues mainly with respect to vertical linear variables. Vertical linear photogrammetric variables of the soft cutaneous tissues could thus be used to evaluate the cephalometric variables of the hard tissues in the vertical direction, when the X-ray is considered too invasive. This helps to significantly reduce the risk of irradiation of subjects when evaluating facial skeletal structures for epidemiological purposes in studies involving a larger number of subjects.

Authors' Contribution: All the authors contributed to the manuscript

Financial Support - Funded by the authors

Conflict of Interest - None to declare

References

1. Jain SK, Anand C, Ghosh SK. Photometric facial analysis-A baseline study. *J. Anat. Soc. India* 2004; 53 (2): 11-13.
2. Wen YF, Wong HM, Lin R, Yin G, McGrath C. Inter-Ethnic / Racial Facial Variations: A Systematic Review and Bayesian Meta-Analysis of Photogrammetric Studies. *PLoS One*. 2015; 10 (8): e0134525.
3. Subtelny JD. A longitudinal study of soft tissue structures and their profile characteristics, defined in relation to underlying skeletal structures. *American Journal of Orthodontics*. 1959; 45 (7): 481-507.
4. Arnett GW, Bergman RT. Facial keys to orthodontic diagnosis and treatment planning. Part I. *Am J Orthod Dentofacial Orthop*. 1993; 103 (4): 299-312.

5. Arnett GW, Bergman RT. Facial keys to orthodontic diagnosis and treatment planning - Part II. *Am J Orthod Dentofacial Orthop*. 1993; 103 (5): 395-411.
6. Bergman RT. Cephalometric soft tissue facial analysis. *Am J Orthod Dentofacial Orthop* 1999; 116 (40): 373-389.
7. Park HS, Rhee SC, Kang SR, Lee JH. Harmonized profiloplasty using balanced angular profile analysis. *Aesthetic Plastic Surgery* 2004; 28: 89-97.
8. Phillips C, Greer J., Vig P., Matteson S. Photocephalometry: errors in projection and landmark location. *Am J Orthod*. 1984; 86 (3): 233-243.
9. Del Santo LM, De Souza RP, Del Santo M JR, Marcantônio E. Alterações no perfil dos lábios de pacientes submetidos a maxilares avulsos em ortodontia do tipo Fort I R Dental Press Ortodon Ortop Facial 2004; 9 (5): 49-63.
10. Ramos AL, Sakima MT, AS Pinto, Bowman SJ. Upper lip changes correlated to maxillary incisor retraction - a metallic implant study. *Orthod Angle* 2005; 75 (4): 499-505.
11. Silvera GA, Correa FA, Vedovello M FILHO, Valdrigh HC, Vedovello SA, Such EZ. Alterations of the nasolabial joint and the incisivo central incisivo superior pós-tratamento ortodôntico. *Ortodontia*. 2006; 39 (1): 31-36.
12. Wen-Ching KO E, Figueroa AA, Polley JW. Soft tissue profile changes after maxillary advancement with distraction osteogenesis by use of rigid external distraction device: a 1- year follow-up. *J Oral Maxillofac Surg* 2000; 58 (9): 959-969.
13. Diouf JS, Diop Ba K, Badiane A, Ndiaye M, Ngom PI, Diagne F. Vertical photogrammetric characteristics of soft tissue with some Senegalese subjects. *Rev. Col. Odonto-Stomatol Afr Chir Maxillo-fac*. 2014; 21 (2): 21-27.
14. Diouf JS, Badiane A, Ngom PI, Niagha G, Diop Ba K, Diagne F. Cephalometric characteristics of subjects received for orthodontic consultation at the clinic of the Department of Odontology of Dakar. *Tropical Odonto- Stomatology* 2010; 33 (3): 27-34.
15. Ferrario VF, Sforza C, Miani A, G. Tartaglia Craniofacial morphometry by photographic evaluations. *American Journal of Orthodontics and Dentofacial Orthopedics*. 1993; 103 (4): 327-337.
16. Cohen J. *Statistical Power and Analysis for Behavioral Science* (2nd ed.). Hillsdale NJ: Lawrence Erlbaum Associates. 1988.
17. Park YC, Burstone CJ. Soft-tissue profile - fallacies of hard-tissue standard in treatment planning. *Am J Orthod Dentofacial Orthop* 1986; 90: 52-62.
18. Carvalho Rosas Gomes L, Horta KOC, LG Gandini, Gonçalves M, Gonçalves JR. Photographic assessment of cephalometric measurements. *Orthod Angle* 2013; 83 (6): 1049-1058.
19. Adwani K, Swami V, Patil A. Correlation between cephalometric and photogrammetric measurements used in orthodontic diagnosis. *Journal of Contemporary Orthodontics*. 2017; 1 (3): 1-4.

Pattern of Malocclusion seen at Aminu Kano Teaching Hospital, Kano Nigeria: A 5 Year Review

*Jibril M, Oguchi C, Adeyemi TE, Yahaya A.

Abstract

Background: Malocclusion involves irregular tooth alignment or occlusion beyond normal limits, influenced by genetic, environmental, and ethnic factors. It is a multifactorial condition without a single cause. Understanding malocclusion patterns across populations is essential for orthodontic treatment planning, especially as demand for corrective care continues to rise globally.

Methods: This retrospective cross-sectional study analysed data from 106 orthodontic patients aged 8 to 40 years seen at Aminu Kano Teaching Hospital (AKTH) from 2019 to 2024. All patients who presented at the Orthodontic clinic of AKTH were included in the study sample. Angles malocclusion types, overjet, overbite, crowding, diastema, and oral habits were assessed. Data was cleaned and processed in MS Excel and all statistical analysis was performed using the Statistical Package for Social Sciences (SPSS) version 20, SPSS Inc., Chicago, IL, USA. (SPSS).

Results: The 5-year study at Aminu Kano Teaching Hospital included (106) patients, predominantly female (65.7%), with an average age of 14.5 years. Most patients (40%) were aged 11-15 years. A large proportion of the subjects (64.4%) had Class I malocclusion, with low diastema prevalence (89.5% at 0-1 mm). Increased overjet affected 51.4% and 50.5% were affected by increased overbite. This shows that a small majority had increased overjet (51.4%) and increased overbite (50.5%), a few 16.2% had crossbite, and 7.6% anterior open bite, while a majority of the subjects 69.2% had tooth rotations. Malocclusion distribution showed no gender differences but varied by age. Class I malocclusion was highest in age groups <10 and >25 years, while anterior open bite was more prevalent in older age groups. Significant associations were found with overbite and anterior open bite. Approximately 31.4% of patients exhibited oral habits, with nail biting being most common. No significant gender differences were observed in the prevalence of oral habits. Oral habits were not significantly associated with malocclusion types except for anterior open bite, where 87.5% of affected patients exhibited habits. Consider the only significant relationship between oral habits and malocclusion type was observed between anterior open bite (AOB) where 87.5% of subjects who had AOB gave a positive history of oral habits.

Conclusion: This study highlights the prevalence of malocclusion, different malocclusion types, low diastema rates, and significant occurrences of increased overjet, overbite, and tooth rotations, emphasising the need for regional assessments to address the dental needs of the population.

Authors' Affiliation

Department of Child Dental Health,
Bayero University, Kano/ Aminu Kano Teaching Hospital,
Kano, Nigeria.

Correspondence

*Dr M Jibril

Department of Child Dental Health
Bayero University, Kano/ Aminu Kano Teaching Hospital,
Kano.

Email address: drjibril2@gmail.com

Phone: +2347037055374

Introduction

Malocclusion refers to an irregularity concerning teeth alignment and/or their relationship during dental occlusion

beyond the range of what is accepted as normal.¹ Malocclusion ranks amongst the top 3 oral pathologies and oral public health priorities.² Numerous etiological factors have been suggested for malocclusion, with genetics, environmental influences, and ethnic background being primary contributors.³ Functional adaptations to environmental factors can impact surrounding structures, such as dentition, bone, and soft tissues, ultimately leading to various malocclusion issues. Consequently, malocclusion is considered a multifactorial condition without a single definitive cause.⁴

Several studies^{4,5,6} on malocclusion have reported varied findings across different populations worldwide.⁷ There is clear evidence of ethnic variation in prevalence and types of malocclusion. As the demand for orthodontic treatment increases in many countries,⁷ with data showing the negative impact of malocclusion on quality of life, understanding the patterns of common malocclusions in specific populations will aid orthodontic practitioners in planning effective treatment.

There is limited data regarding the pattern of malocclusion in Northern Nigeria. The aim of this study was to determine the types of malocclusion based on gender and age among orthodontic patients at Aminu Kano Teaching Hospital (AKTH), Kano, and to determine the pattern of distribution of the anterior posterior relationships of the jaws and the effects of oral habits on these malocclusion patterns. Furthermore, the data will be valuable for comparing the results of this study with findings reported in other populations.

Materials and methods

This descriptive retrospective study analysed data collected from pre-treatment records of patients seen between January 2019 to July 2024 at the Orthodontic Unit of the Child Dental Health Department, Aminu Kano Teaching Hospital (AKTH) in Kano. Records of a total of 106 patients aged 8 to 40 years were reviewed and collected.

The inclusion criteria for the study were as follows:

1. Patients who presented at the Orthodontic Unit of between January 2019 and July 2024 with compliant of malocclusion or related symptoms
2. Subjects who had complete pre-treatment records
3. Presence of first permanent molars and canines
4. No prior history of orthodontic treatment

The exclusion criteria were:

1. Patients with incomplete orthodontic records
2. Patients without first permanent molars or canines
3. Patients with significantly deteriorated first permanent molars

Demographic data (age and gender) and malocclusion type were recorded for each patient. Molar relationships were classified using Angle's classification of malocclusion into Class I, Class II, or Class III malocclusion. Overjet was measured in millimetres from the edge of the upper central incisor to the labial surface of the lower central incisor; 1–3 mm was considered normal, greater than 3 mm was classified as increased, and less than 1 mm was reduced. Overbite was also measured in millimetres as the perpendicular distance from the edge of the lower central incisor to the upper central incisor. A normal range was 0–3 mm; values above 3 mm indicated a deep bite, and values below 0 mm indicated an open bite.

Crowding in the upper and lower arches was measured in millimetres and categorised as follows: 0 mm (no crowding), 1–3 mm (mild crowding), 4–6 mm (moderate crowding), and 7 mm and above as severe crowding. A maxillary midline diastema was diagnosed when there was a space of at least 1 mm between the upper central incisors. Oral habits were noted as present or absent, with specific types documented.

Ethical clearance for the study was obtained from the institutional Health research and Ethics board of Aminu Kano Teaching Hospital.

Data was cleaned and processed in MS Excel and statistical analysis was performed using the Statistical Package for Social Sciences (SPSS) version 20, SPSS Inc., Chicago, IL, USA. (SPSS).

Results

Socio-demographic data of the study participants

The orthodontic patient population at the Aminu Kano Teaching Hospital, Kano, Nigeria, over the 5-year period from 2019 to 2024, comprised 106 patients. Female patients outnumbered their male counterparts, accounting for 65.7% of the total patient population (Table 1). The age range of these patients spanned from 8 to 40 years, with a mean age of 14.5 ± 5.6 years. The largest proportion of patients (40%) fell within the 11–15 years age bracket. This

was followed by the <10 years age group, which accounted for 26.7% of the patients. The remaining age groups had smaller proportions, with the oldest

age group (>25 years) comprising only 5.7% of the patients.

Table 1. Age and sex distribution of orthodontic patients at the Aminu Kano Teaching Hospital, Kano (2019 – 2024)

Sex	n (%)	Range	Mean $\pm$ SD	P-value
Female	69 (65.7)	7–40	14.7 $\pm$ 5.6	0.72
Male	36 (34.3)	8–32	14.3 $\pm$ 5.6	
Total	105 (100)	7–40	14.5 $\pm$ 5.6	

P-value estimated using Independent t-test

A comparison of age group distribution by sex (Figure 1) indicates that female patients were more prevalent in the younger age groups, while the proportions of male and female patients were more

evenly distributed in the older age groups. However, there was no statistically significant difference in the mean age between male and female patients ($p = 0.72$).




Figure 1. Age group distribution by sex among orthodontic patients at the Aminu Kano Teaching Hospital, Kano (2019 – 2024)

Overall prevalence and distribution pattern
Table 2 summarises the pattern of malocclusion among orthodontic patients treated at the Aminu Kano Teaching Hospital, Kano, Nigeria, between

2019 and 2024. Based on Angle's Classification of Malocclusion, the majority of patients (64.4%) exhibited Class I malocclusion, followed by Class II (29.8%) and Class III (5.8%) malocclusions.

The prevalence of diastema, defined as spacing between the teeth of more than 1mm was low, with 89.5% of patients having a diastema of 0 to 1 mm. Overjet was recorded in 51.4% of patients, with 44.8% having an overjet of 0 to 4 mm. Regarding overbite, 50.5% of patients had an overbite of 2 to 3

mm, while 45.7% exhibited an overbite of 4 mm or more. Crossbite was present in 16.2% of patients. Additional findings revealed that 7.6% of patients had an anterior open bite, and 69.2% showed tooth rotations (Table 2).

Table 2. Pattern of malocclusion among orthodontic patients at the Aminu Kano Teaching Hospital, Kano (2019 – 2024)

Malocclusion	Frequency, n	Percentage, %	95% CI
Angle's Class			
Class I	67	64.4	54.8 – 73.0
Class II	31	29.8	21.0 – 39.2
Diastema			
<1 mm	94	89.5	82.0 – 94.6
>1 mm	11	10.5	5.3 – 18.0
Overjet			
0 – 4 mm	47	44.8	35.0 – 54.8
> 4 mm	54	51.4	41.5 – 61.3
Reverse	4	3.8	1.0 – 9.5
Overbite			
0 – 1 mm	4	3.8	1.0 – 9.5
2 – 3 mm	53	50.5	40.5 – 60.4
≥ 4 mm	48	45.7	35.9 – 55.7
Crossbite			
Absent	88	83.8	75.3 – 90.3
Present	17	16.2	9.7 – 24.6
Anterior Open Bite			
Absent	97	92.4	85.5 – 96.7
Present	8	7.6	3.5 – 14.5
Rotations			
Absent	32	30.8	21.9 – 40.2
Present	72	69.2	58.8 – 77.3

Distribution of malocclusion according to gender and age

The distribution of malocclusion in the study population was analysed by sex and age groups (Tables 3 and 4). There was no statistically significant difference in the distribution of Angle's

Classification, diastema, overjet, overbite, crossbite, anterior open bite, and rotations between male and female patients.

In contrast, analysis by age groups revealed notable differences. The highest proportion of Angle's Class I malocclusion was observed in the <10 years (75%)

and >25 years (83.3%) age groups. Conversely, Class III malocclusion was more prevalent in the 16–20 and 21–25 age groups (17.4% and 16.7%, respectively, $p = 0.06$).

Additionally, younger age groups (<10 years and 11–15 years) had a significantly higher proportion of increased overbite (≥ 4 mm) compared to older age

groups ($p = 0.003$). The analysis also revealed a significant association between age and anterior open bite ($p = 0.05$), with the prevalence of anterior open bite being highest among older age groups: 33.3% in patients aged 21–25 years and 16.7% in those above 25 years.

Table 3. Pattern of Malocclusion according to gender and age among orthodontic patients at the Aminu Kano Teaching Hospital, Kano (2019 – 2024)

Malocclusion	Sex, n (%)		Age (years), n (%)				
	Female (n = 68).	Male (n = 36)	<10 (n = 28)	11 – 15 (n = 42)	16 – 20 (n = 23)	21 – 25 (n = 6)	>25 (n = 6)
Angle's Class							
Class I	43(63.2).	24 (66.7)	21(75.0)	28(68.3)	11(47.8)	2(33.3)	5(83.3)
Class II.	21(30.9).	10 (21.8)	6(21.4)	13(31.7)	8(34.8)	3(50.0)	1(16.7)
Class III	4(5.9)	2 (5.6)	1(3.6)	0	4(17.4)	1(16.7)	0
<i>p</i> = 0.94		<i>p</i> = 0.06					
Diastema							
<1mm	63(91.3)	31(86.1)	24(58.7)	37(88.1)	23(100)	5(83.3)	5(83.3)
>1mm	6(8.7)	5(13.9)	4(14.3)	5(11.9)	0	1(16.7)	1(16.7)
<i>p</i> = 0.41		<i>p</i> = 0.23					
Overjet							
0 – 4 mm	30 (43.5)	17(47.2)	15(53.6)	16(38.1)	9(39.1)	4(66.7)	3(50.0)
> 4 mm.	37(53.6)	17(47.2)	12(42.9)	25(59.5)	12(52.2)	2(33.3)	3(50.0)
Reverse	2 (2.9)	2(5.6)	1(3.6)	1(2.4)	2(8.7)	0	0
<i>p</i> = 0.66		<i>p</i> = 0.73					

P-value estimated using Chi-square test and Fisher's exact test

Table 4. (Continuation) Pattern of malocclusion according gender and age among orthodontic patients at the Aminu Kano Teaching Hospital, Kano (2019 – 2024)

Malocclusion	Sex, n (%)		Age (years), n (%)				
	Female (n = 68).	Male (n = 36)	<10 (n = 28)	11 – 15 (n = 42)	16 – 20 (n = 23)	21 – 25 (n = 6)	>25 (n = 6)
Overbite							
0 – 1 mm	2(2.9)	2(5.6)	0	0	4(17.4)	0	0
2 – 3 mm	33(47.8)	20(55.6)	19(67.9)	14(33.3)	12(52.2)	4(66.7)	4(66.7)
≥ 4 mm	34(49.3)	14(38.9)	9(32.1)	28(66.7)	7(30.4)	2(33.3)	2(33.3)
	$p = 0.50$		$p = 0.003$				

Table 4. (Continuation) Pattern of malocclusion according gender and age among orthodontic Patients at the Aminu Kano Teaching Hospital, Kano (2019 – 2024) (contd)

Malocclusion	Sex, n (%)		Age (years), n (%)					
	Female (n = 68).	Male (n = 36)	<10 (n = 28)	11 – 15 (n = 42)	16 – 20 (n = 23)	21 – 25 (n = 6)	>25 (n = 6)	
Crossbite								
Absent	58 (84.1)	30 (83.3)	22 (78.6)	37 (88.1)	18 (78.3)	6 (100)	5 (83.3)	
Present	11 (15.9)	6 (16.7)	6 (21.4)	5 (11.9)	5 (21.7)	0	1 (16.7)	
		<i>p</i> = 0.92		<i>p</i> = 0.62				
Anterior Open Bite								
Absent	64 (92.7)	33 (91.7)	27 (96.4)	38 (90.5)	23 (100)	4 (66.7)	5 (83.3)	
Present	5 (7.2)	3 (8.3)	1 (3.6)	4 (9.5)	0	(33.3)	1 (16.7)	
		<i>p</i> = 1.00		<i>p</i> = 0.05				
Rotations								
Absent	23 (33.3)	9 (25.7)	10 (37.0)	11 (26.2)	8 (34.8)	2 (33.3)	1 (16.7)	
Present	46 (66.7)	26 (74.3)	17 (63.0)	31 (73.8)	15 (65.2)	4 (66.7)	5 (83.3)	
		<i>p</i> = 0.43		<i>p</i> = 0.81				
P-value estimated using Chi-square test and Fisher's exact test								

Prevalence of Oral Habits

Table 5 presents the prevalence and distribution of oral habits among the study population. Approximately one-third of the patients (31.4%) exhibited at least one oral habit. The most common habit was nail biting, observed in 33.3% of patients with oral habits (Figure 2). Tongue thrusting and lip biting were also frequent, each affecting 18.2% of

patients. Other habits, such as thumb sucking, mouth breathing, and teeth gnashing, were less prevalent.

When analysed by gender, oral habits were found in 27.5% of female patients and 38.9% of male patients. However, this difference was not statistically significant ($p = 0.23$), indicating a comparable prevalence of oral habits between male and female patients.

Table 5. Prevalence and distribution of oral habits by gender

Gender	N	Oral Habit, n (%)		95% CI of	P-value
		Absent	Present	Prevalence	
Female	69	50 (72.5)	19 (27.5)	17.5 – 39.6	0.23
Male	36	22 (61.1)	14 (38.9)	23.1 – 56.5	
Overall	105	72 (68.6)	33 (31.4)	22.7 – 41.2	
P-value estimated using Chi-square test					




Figure 2. Type of oral habit among orthodontic patients at the Aminu Kano Teaching Hospital, Kano (2019 – 2024)

Association Between the Presence of Oral Habits and Malocclusion

Table 6 illustrates the association between oral habits and malocclusion. The findings indicate that oral habits were not significantly associated with Angle's Classification ($p = 0.37$), overjet ($p = 0.21$), overbite ($p = 0.21$), or crossbite ($p = 0.44$).

However, a marginally significant association was observed between oral habits and diastema ($p = 0.10$).

Over half (54.5%) of patients with a larger diastema (2–4 mm) exhibited an oral habit, compared to 28.7% of patients with normal spacing (0–1 mm).

The only significant association identified was between oral habits and anterior open bite ($p < 0.001$). A striking 87.5% of patients with anterior open bite had an oral habit, compared to 26.8% of those without anterior open bite.

Table 6. Association between the presence of oral habits and malocclusion among orthodontic Patients at the Aminu Kano Teaching Hospital, Kano (2019 – 2024)

Malocclusion	N	Oral Habit, n (%)		P-value
		Absent	Present	
Angle's Class				
Class I	67	43 (64.2)	24 (35.8)	0.37
Class II	31	24 (77.4)	7 (22.6)	
Class III	6	5 (83.3)	1 (16.7)	
Diastema				
< 1mm	94	67 (71.3)	27 (28.7)	0.10
> 1mm	11	5 (45.4)	6 (54.5)	

Overjet				
0 – 4 mm	47	28 (59.6)	19 (40.4)	0.21
> 4 mm	54	41 (75.9)	13 (24.1)	
Reverse	4	3 (75.1)	1 (25.0)	
Overbite				
0 – 1 mm	4	3 (75.0)	1 (25.0)	0.21
2 – 3 mm	53	32 (60.4)	21 (39.6)	
≥ 4 mm	48	37 (77.1)	11 (22.9)	
Crossbite				
Absent	88	59 (67.0)	29 (32.9)	0.44
Present	17	13 (76.5)	4 (23.5)	
Anterior Open Bite				
Absent	97	71 (73.2)	26 (26.8)	< 0.001
Present	8	1 (12.5)	7 (87.5)	
Rotations				
Absent	32	22 (68.7)	10 (31.2)	0.94
Present	72	50 (69.4)	22 (30.6)	

P-value estimated using Chi-square test and Fisher's exact test

Discussion

The pattern of malocclusion varies globally, regionally, and even between cities.⁸ Evaluating orthodontic patients provides valuable insights for planning treatment and assessing malocclusion distribution. Understanding these patterns is essential for guiding treatment priorities and developing effective orthodontic services and preventive programs tailored to specific populations.⁹

The majority of participants in this study were female, a finding consistent with previous studies,^{10,11} indicating that women are more likely to seek orthodontic treatment. This trend can be attributed to the greater emphasis that females typically place on dental aesthetics and appearance, compared to males. Social and cultural factors may also play a role, as women are often subjected to societal expectations regarding physical appearance, which can influence their health-seeking behaviours, including pursuing treatments to enhance their smiles. Additionally, studies^{12,13} have suggested that females may have a

higher level of self-awareness and concern about oral health, contributing to their proactive approach toward orthodontic care.

Angle's Class I malocclusion was the most prevalent pattern observed in 64.4% of participants who sought treatment at the Orthodontic Unit of the Child Dental Health Department, Aminu Kano Teaching Hospital. This finding is consistent with studies conducted in the southern regions of Nigeria,^{10,14-16} other African countries,^{3,17} and globally.^{1,5} The global prevalence of Angles class 1 Malocclusion could be related to the wide range of malocclusion traits within this class.

In this study, Angle's Class II malocclusion was observed in 29.8% of participants, a value comparable to the findings reported by Borzabadi-Farahani *et al.*¹⁸ among urban Iranian adolescents but lower than those reported by Guclipaneni *et al.*¹⁹ among Saudi adolescents. Class II malocclusion has been found to be the most common pattern among individuals of Northern European descent²⁰ and Pakistani²¹ populations. The proportion of subjects

with Angles Class II malocclusion in our study was higher than reported in some studies from Southern Nigeria,^{6,14,22} but similar to results from another study in Lagos, Nigeria.¹¹ Angle's Class III malocclusion was the least common in this study, found in 5.8% of participants. This is higher than the findings reported by Ajayi²² (1.8%) among schoolchildren in Benin City and Obanubi *et al.*¹¹ (4.1%) among patients at the Lagos State University Teaching Hospital. However, it is lower than findings by Onyeaso *et al.*⁶ (8.0%) at the University College Hospital Ibadan and Folaranmi and Okeke.¹⁰ (24.3%) at the University of Nigeria Teaching Hospital, Enugu. The variations in malocclusion patterns across studies may be attributed to differences in sample selection techniques, the diversity of local ethnic groups, and cultural practices. Overall the relatively low frequency of Class III malocclusion aligns with craniofacial norms reported in sub Saharan Africa.

Overall, the findings from this study is similar with the studies reported in other parts of Nigeria,^{9,10} Africa,^{3,17} and globally,^{1,5} where Class I malocclusion remains the most common, followed by Class II and Class III malocclusions. These results provide valuable insights for orthodontic practitioners to develop targeted treatment strategies tailored to the specific needs of different populations.

In the present study, 74.3% of participants had Skeletal Class I pattern, indicating a normal skeletal relationship between the upper and lower jaws. Skeletal Class II and Class III patterns were observed in 19.0% and 6.7% of patients, respectively.

The present study also found a low prevalence of diastema, (space greater than 1mm) in 10.5%, contrasting with higher rates reported in studies from the South-Southern²² and South-Western⁶ regions of Nigeria. The variations noticed in diastema prevalence may be attributed to differences in demographics, age groups, and cultural or genetic factors. Our study recorded high values of both overjet and overbite, with overjet being more prevalent than overbite. These findings differ from studies conducted in the southern regions of Nigeria

and among Tanzanian children,²³ where higher occurrences of normal overjet and overbite were reported. The variations in findings may be partially attributed to differences in the criteria used to define increased overjet in various studies. Discrepancies in sample selection, measurement techniques, and population demographics could also contribute to these differences. In this study, crossbite was observed in 16.2% of participants, a prevalence higher than the values reported by Ajayi²² (11.5%) and Onyeaso *et al.*⁶ (12.8%). However, it was lower compared to the findings by Otuyemi *et al.*²⁴ (17.8%) and Utomi *et al.*²⁵ (20.4%).

Furthermore, the study revealed that 7.6% of patients had an anterior open bite, a value comparable to the 7.1% reported by Aikins and Onyeaso in Rivers State.¹⁵ However, it was higher than the values reported by Ajayi²¹ (4.1%) in Benin and Gudipaneni *et al.*¹⁹ (4.6%) among Saudi subjects. Anterior open bite is clinically significant due to its impact on aesthetics, speech, and biting efficiency. Additionally, 69.2% of the participants exhibited tooth rotations. The high prevalence of tooth rotations emphasises the need for corrective interventions to improve occlusion, dental hygiene, and overall smile aesthetics.

This study found no significant association between malocclusion patterns and participants' age or gender, consistent with findings from Obanubi *et al.*¹¹ and Kashif *et al.*²⁶ However, a higher prevalence (75%) of Angle's Class I malocclusion was observed among participants under 10 years compared to older age groups. This difference may be attributed to parental influence or peer pressure, as younger children are more likely to receive treatment at the recommendation of their parents or caregivers.

These results highlight the importance of regional assessments to inform tailored orthodontic treatment strategies. In addition, given the limited resources for healthcare provision, tailored preventive strategies may be developed along these identified areas. These differences could be attributed to differences in sample size, age distribution, regional factors, and

diagnostic criteria used in each study. Environmental factors, dietary habits, and genetic variations could also play a role in influencing the development of crossbite in different populations.

Conclusion

This study provides important insights into malocclusion patterns among orthodontic patients at Aminu Kano Teaching Hospital, Kano, Nigeria. Angle's Class I malocclusion was the most prevalent, followed by Class II and Class III, aligning with regional and global trends. Diastema was uncommon, but there were notable occurrences of overjet, overbite, crossbite, and tooth rotations, emphasising the need for corrective interventions.

References

1. Alhammadi MS, Halboub E, Fayed MS, Labib A, El-Saaidi C. Global distribution of malocclusion traits: A systematic review. *Dental Press J Orthod* 2018;23:40.e1-40.e10
2. Peres MA, Macpherson LM, Weyant RJ, Daly B, Venturelli R, Mathur MR, Listl S, Celeste RK, Guarnizo-Herreño CC, Kearns C, Benzian H. Oral diseases: a global public health challenge. *The Lancet*. 2019 Jul 20;394(10194):249-60.
3. Pobe R.D , Amoah G.K , Newman-Nartey M , Ndanu T.A , Mayari Yabang G.T. Patterns and distribution of malocclusion among Ghanaian orthodontic population. *GDJ*. 2024; 21(1):4-7.
4. Celikoglu M, Akpınar S, Yavuz I. The pattern of malocclusion in a sample of orthodontic patients from Turkey. *Med Oral Patol Oral Cir Bucal*. 2010; 1:15(5):e791-796
5. Ciuffolo F, Manzoli L, D'Attilio M, Tecco S, Muratore F, Festa F, et al. Prevalence and distribution by gender of occlusal characteristics in a sample of Italian secondary school students: a cross-sectional study. *Eur J Orthod*. 2005;27:601-606.
6. Onyeaso CO, Aderinokun GA, Arowojolu MO. The pattern of malocclusion among orthodontic patients seen in Dental Centre, University College Hospital, Ibadan, Nigeria. *Afr J Med Med Sci*. 2002;31:207-211.
7. Rahman MM, Jahan H, Hossain MZ. Pattern of malocclusion in patients seeking orthodontic treatment at Dhaka Dental College and Hospital. *Ban J Orthod and Dentofacial Orthop*. 2015; 3:9-11.
8. Lone IM, Midlej K, Zohud O, Paddenber E, Krohn S, et al. Global Map of Skeletal and Dental Malocclusion Prevalence: From Classes to Continents. *J Dent & Oral Disord*. 2024; 10(1): 1183.
9. Lin M, Xie C, Yang H, Wu C, Ren A. Prevalence of malocclusion in Chinese schoolchildren from 1991 to 2018: A systematic review and meta-analysis. *Int J Paediatr Dent*. 2020; 30: 144–155.
10. Folaranmi N, Okeke A. A retrospective evaluation of the class of malocclusion amongst orthodontic patients at the university of Nigeria teaching hospital (UNTH), Enugu, Nigeria. *Ann Med Health Sci Res*. 2011;1(1):103-106.
11. Obanubi KO, Ogunbanjo BO, Adeniyi AA and Adegbite AA. The pattern of malocclusion at the orthodontic unit, Lagos state Uni. teaching hospital. *Nig J Clin Med* 2009;1(2): 8-10.

12. Nugroho MJ, Ismah N, Purbiati M. Orthodontic treatment need assessed by malocclusion severity using the dental health component of i o t n . J I n t D e n t M e d R e s . 2019;12(3):1042–1046.
13. Tshepiso DN, Thomas KM, Millicent M. et al. The distribution of malocclusion using the index of orthodontic treatment needs at a university dental hospital in and around Pretoria, South Africa. *International Journal of Oral Health Dentistry* 2023;9(4):258–258.
14. dacosta O, and Utomi I. Referral mode and pattern of malocclusion among patients attending the Lagos university teaching hospital, Lagos, Nigeria. *OST-TDJ* 2009;32:(4):17-23.
15. Aikins EA and Onyiaso CO. Prevalence of malocclusion and occlusal traits among adolescents and young adults in Rivers State, Nigeria. *Odontostomatol Trop*, 2014; 37(145):5-12.
16. Yemitan TA, Afolabi OO. Prevalence of malocclusion in Africa: A systematic review and meta-analysis. *Magna Scientia Advanced Research and Reviews* 2022; 5(01):030-035.
17. Mtaya M, Brudvik P, Astrøm AN. Prevalence of malocclusion and its relationship with socio-demographic factors, dental caries, and oral hygiene in 12- to 14-year-old Tanzanian schoolchildren. *Eur J Orthod*. 2009;31(5):467-476.
18. Borzabadi-Farahani A, Borzabadi-Farahani A, Eslamipour F. Malocclusion and occlusal traits in an urban Iranian population. An epidemiological study of 11- to 14-year-old children. *Eur J Orthod*. 2009;31(5):477–484.
19. Gudipani RK, Aldahmeshi RF, Patil SR, Alam MK. The prevalence of malocclusion and the need for orthodontic treatment among adolescents in the northern border region of Saudi Arabia: an epidemiological study. *BMC Oral Health*. 2018;18(1):16
20. Lone IM, Zohud O, Midlej K, Proff P, Watted N, Iraqi FA. Skeletal Class II Malocclusion: From Clinical Treatment Strategies to the Roadmap in Identifying the Genetic Bases of Development in Humans with the Support of the Collaborative Cross Mouse Population. *J Clin Med*. 2023;12(15):5148.
21. Gul-e-Erum, Fida M. Pattern of malocclusion in orthodontic patients: a hospital based study. *J Ayub Med Coll Abbottabad*. 2008;20(1):43-47.
22. Ajayi EO. Prevalence of Malocclusion among school children in Benin city, Nigeria. *J Biomed Res*. 2008;(2):58–65.
23. Mtaya M, Brudvik P, Astrøm AN. Prevalence of malocclusion and its relationship with socio-demographic factors, dental caries, and oral hygiene in 12- to 14-year-old Tanzanian schoolchildren. *Eur J Orthod*. 2009;31(5):467-476.
24. Otuyemi OD, Isiekwe MC, Sote EO, Jones S. Need for interceptive orthodontic treatment in 3-5 year old Nigerian children. *Paediatr Dent J* 1997; 7:7-11.
25. Utomi IL, Agbonikhena AJ, Isiekwe MC, daCosta OD, Sanu OO. Preventive and interceptive orthodontic treatment need in a Nigerian teaching hospital. *Nig Dent J* 2024; 22:77-81.
26. Aslam K, Nadim R, Rizwan S. Prevalence of angles of malocclusion according to age groups and gender. *Pak Oral Dent J* 2014;34(2):362-365.

The Role of Artificial Intelligence in Orthodontics

*Awwal NM, Aladejare AL

Abstract

Artificial Intelligence (AI) has rapidly emerged as a transformative tool in orthodontics, enhancing diagnostic accuracy, treatment planning, and clinical workflows. From its early conceptualisation to its present integration into healthcare, AI now plays a pivotal role in optimising orthodontic outcomes through data-driven decision-making and automation. This review synthesises current literature on the applications, benefits, and limitations of AI in orthodontics. It explores AI paradigms, including symbolic AI, machine learning, and deep learning, while categorising their functionalities into classification, regression, detection, and segmentation. Key databases and recent studies were evaluated to gather relevant data and outcomes across multiple subdomains of orthodontic practice. AI has been effectively implemented in diverse orthodontic tasks such as cephalometric analysis, bone age prediction, airway assessment, facial proportion analysis, and appliance fabrication. AI-assisted systems have demonstrated high accuracy (often >90%) in diagnosis, treatment planning, and prediction models. Notably, applications like extraction decision support, impacted canine management, and orthognathic surgery planning have shown significant promise. The integration of AI with teleorthodontics and 3D printing also opens new avenues for remote and customised care. AI is reshaping orthodontic practice by improving precision, efficiency, and patient outcomes. While challenges such as data bias, privacy concerns, and regulatory gaps remain, ongoing innovations suggest a future where AI could democratise access to advanced orthodontic care, especially in underserved regions. Strategic implementation and ethical governance will be key to its successful integration into routine practice.

Authors' Affiliation

Orthodontic unit,
Department of Child Dental Health
Lagos University Teaching Hospital, Lagos, Nigeria.

Correspondence

*Dr NM Awwal
Orthodontic unit,
Department of Child Dental Health
Lagos University Teaching Hospital, Lagos, Nigeria.
Email address:
Phone number: +2348067645220

Introduction

Artificial Intelligence (AI), a term first introduced in 1955 by John McCarthy, describes the ability of machines to perform tasks that are classified as intelligent.^{1,2} It is the simulation of human intelligence in machines, enabling them to learn, reason, and make decisions with minimal human input.²⁻⁸ AI systems can analyse vast amounts of data, recognise patterns, and make

decisions with minimal human intervention, improving efficiency and automation in various fields.³ In orthodontics, AI enhances diagnostic accuracy, treatment planning, and overall patient care by analysing large datasets, recognising patterns, and predicting outcomes with minimal human intervention.⁸

Paradigms of AI in Orthodontics

1. Symbolic AI (GOFAI): Symbolic AI, also referred to as "good old-fashioned AI" (GOFAI) which was popular until the late 1980s, uses rules, such that when a certain criterion is met, a corresponding action must be taken.⁶ Rule-based systems using "if-then" logic.⁶
2. Machine Learning (ML)⁴: Machine learning is a term first phrased by Arthur Samuel in 1952.⁵ In machine learning (ML), algorithms are classified according to the kind of learning and the desired

result. There are three main categories: ⁴
Supervised learning: Supervised learning is a procedure used in orthodontic decision-making, such as extraction planning, where predictions or classifications are guided by known outcomes.⁴ In supervised learning the output data is known in advance and this helps AI to be able to make precise predictions about a newly input data.⁸

Unsupervised learning: In orthodontics, unsupervised learning arranges data without labels, exposing patterns or connections. It helps with tasks that do not have defined results, making analyses and classifications easier. ^{4,8,13}

Reinforced learning: Reinforced learning helps systems improve by learning through trial and error using feedback, without prior knowledge. In orthodontics, this approach enhances efficiency by

enabling systems to learn from mistakes and refine their methods over time.⁴

3. Deep Learning (DL): Deep learning (DL) is a branch of machine learning that automatically extracts features from raw data. It uses artificial neural networks (ANNs), developed in the 1990s.⁵ With better computing power, deeper and more complex networks have been created. These “deep” networks can solve more advanced problems. DL reduces the need for manual input. It learns patterns from data on its own. This makes it useful in areas like image analysis and healthcare.⁵ As a subset of machine learning, deep learning uses neural networks and has excelled in processing complex data structures, particularly high-dimensional data such as images.^{7,10}




Figure 1: Shows a simplified AI diagram, which shows machine learning as a paradigm of artificial intelligence and deep learning, a subset of machine learning.

Four AI driven tasks

The 4 major AI-driven tasks in dentistry are :

1. Classification
2. Regression
3. Detection, and
4. Segmentation

(1) Classification is the most known AI-driven task and it assigns objects to categories that have been pre-specified or pre-labeled.⁹ Examples include; support

vector machine (SVM), decision trees and many more.⁹ (2) Regression analysis is a statistical and machine learning method used to identify relationships between variables and predict continuous outcomes. For example, it can estimate a patient's age based on tooth wear or bone density by finding patterns in the data.⁹ (3) Object detection focuses on determining the location of objects within

an image or video. Unlike basic image classification, which merely identifies the presence of objects, detection goes a step further by pinpointing their exact positions. A widely used method for object detection is the Region-based Convolutional Neural Network (R-CNN), which involves generating region proposals, extracting features using a CNN, and classifying each proposed region.⁹ (4) In

dentistry, object segmentation involves pixel-level classification to distinguish different structures within a region of interest.⁹ This technique helps divide dental images into components like teeth, gums, bones, and lesions, aiding diagnosis and treatment planning. In orthodontics, for instance, it allows individual teeth to be isolated for alignment assessment and appliance design.




Figure 2: Key Applications of AI in Orthodontics

Diagnosis and Treatment Planning

The area of orthodontic diagnosis and treatment planning is changing as a result of the integration of artificial intelligence which gives better accuracy and efficiency.⁴ Machine learning and computer vision have made it easy for AI to automate complex patient evaluations and this gives orthodontists access to a wide range of cephalometric analytic tools.⁴ A study by Ezhov (2021) compared AI-assisted and unaided dentists in oral CBCT evaluations. The AI system improved diagnostic performance, with higher sensitivity (0.8537 vs. 0.7672) and specificity (0.9672 vs. 0.9616) in the AI-aided group.⁶

Systematic reviews and meta-analyses on AI for detecting caries and periapical lesions have shown varying accuracy (68%–99.2%) depending on imaging methods.⁶ A 2019 meta-analysis reported 88.75% accuracy for AI in detecting radiolucent lesions, while a 2023 study by Sadr found AI had high sensitivity (0.925) and specificity (0.852).⁶ These findings suggest AI can serve as an effective diagnostic aid in diagnosis and treatment planning.⁶ Due to the ability of AI, machine language, computer vision and natural language processing combined simplify patient diagnosis and treatment planning.⁴




Figure 3: It shows AI depends diagnosis and treatment planning workflow

Bone Age Prediction

The choice of proper treatment timing depends on bone age prediction based on cervical vertebrae maturity, and this is very vital because it directly impacts the effectiveness of orthodontic treatments.⁶ The Machine Learning-based methods, which include the convolutional neural networks, according to Seo et al. (2021), have shown >90% accuracy for vertical and sagittal skeletal maturation diagnosis.⁶ In orthodontics, classification models have shown greater accuracy than clustering methods for predicting bone age. Among various classifiers, the artificial neural network (ANN) achieved the highest performance, with a weighted kappa of 0.926, indicating its effectiveness in determining cervical vertebral maturation.¹⁰

Airway Obstruction Assessment

There is a reciprocal relationship between skeletal deformities and airway obstruction. Impaired airflow, such as that caused by upper-airway obstruction, can influence breathing patterns and disrupt normal craniofacial growth, potentially resulting in malocclusion and other facial irregularities. Therefore, identifying conditions like adenoid hypertrophy is essential for accurate

orthodontic assessment and effective treatment planning.¹¹ In light of this, Dong et al. introduced two deep learning models—hierarchical masks self-attention U-net (HMSAU-Net) for upper airway segmentation and 3D-ResNet for diagnosing adenoid hypertrophy from CBCT scans. The 3D-ResNet10 model demonstrated a high accuracy of 0.912 in diagnosing adenoid hypertrophy.¹²

Landmark Identification and Cephalometry Tracing

The use of artificial intelligence (AI) in identifying anatomical landmarks on radiographs has enhanced orthodontic diagnosis and treatment planning.⁴ In order to advance cephalometric analysis, Montúfar et al. (2018) introduced a hybrid approach that automates the annotation of anatomical landmarks in cephalometric images.¹⁸ This method integrates artificial intelligence (AI) algorithms with dynamic shape models, allowing for more accurate and efficient identification of key landmarks, thus streamlining the process of cephalometric analysis.⁴ In addition, Montúfar et al. (2018) developed an AI-driven system for automatic 3D cephalometric landmark identification, aimed at simplifying the analysis of craniofacial changes in response to orthodontic therapy.¹⁸




Figure 4: Shows a sample of automatic cephalometric landmark tracings performed using WebCeph. It shows that the results of Downs cephalometric analysis are superimposed on the tracing and measurements outside the standard range marked in red with asterisks.

Growth and Development Estimation

Evaluating growth and development indicators is vital for properly timing orthodontic treatment. Key markers like skeletal maturity, physical height, and chronological age play a significant role in orthodontic diagnosis and planning.⁴ K  k et al. (2019) reported that among seven different AI approaches used to evaluate cervical vertebral

maturation, Artificial Neural Networks (ANN) demonstrated the highest performance.¹⁷ By mimicking the structure and function of the human nervous system, Artificial Intelligence (AI) significantly enhances both the precision and speed of analyzing growth and skeletal maturity indicators in orthodontics.⁴




Figure 4: It shows an artificial intelligence-based algorithm for cervical vertebrae maturation stage assessment where AI algorithms are used to automatically identify and outline cervical vertebrae (C2–C4) on lateral cephalometric radiographs. By analyzing the shape and size of these bones, the AI accurately determines the patient's skeletal maturation stage, helping orthodontists plan growth-related treatments more objectively and efficiently.

Facial Proportion Analysis

AI is revolutionising facial proportion assessment by using optical facial recognition, in contrast to traditional method relying on profile photographs and lateral cephalometric radiographs.¹⁹ By surpassing the limitations of traditional dimensional analysis, artificial intelligence (AI) enables the assessment of angular measurements and facial ratios. AI offers objective evaluations of facial aesthetics by incorporating various beauty parameters, potentially enhancing orthodontic and surgical decision-making through a deeper understanding of facial attractiveness.⁴

Extraction Prediction:

Xie et al. created an expert system (ES) using artificial neural networks (ANN) that achieved 80% accuracy. This was done in order to determine whether extraction is needed for management of malocclusion in patients between the ages of 11 and 15.²¹ Jung et al. used artificial neural networks (ANN) to forecast extraction patterns with 84% accuracy.²⁰ Artificial intelligence (AI) technologies simulate human decision-making, achieving 80-90% accuracy in orthodontic extraction decisions. These technologies particularly assist orthodontists with less experience, offering guidance that enhances decision-making. AI algorithms not only optimise treatment outcomes but also reduce human error, streamline processes, and help tailor orthodontic interventions to individual patient needs, improving overall efficiency in clinical practice.⁴

Management of Impacted Canines

AI-assisted Radiographic Analysis: Deep learning models analyse panoramic and CBCT (Cone Beam Computed Tomography) images to detect impacted canines early, even before they become clinically evident.²² A study by Kumar et al. demonstrated that CNN successfully detects impacted and non impacted maxillary canines on cropped and uncropped Panoramic radiographs.²²

AI helps in identifying dental and skeletal landmarks, improving accuracy in localizing impacted canines.

AI models simulate the likely eruption path of impacted canines. It involves the utilisation of CBCT to enhance early diagnosis and interception of canine impactions and this addresses the limitations found in traditional 2D radiographs.²³ Virtual Surgical Planning (VSP): AI assists in planning surgical exposure of impacted canines, optimizing flap design and minimizing complications.²⁴ AI significantly improves precision, speed, and reliability in the management of impacted canines in orthodontics. It aids in early identification, treatment strategy, and follow-up care, leading to fewer mistakes and better patient results.²⁵

Treatment Outcome Assessment

Traditional multi-regression models analyse dental treatment outcomes but are limited by their linear assumptions and inability to capture all possible results.⁴ Artificial neural networks overcome these limitations by modeling complex, non-linear relationships, especially in orthodontic cases like Class II and III malocclusions, further enhanced by the digital clear aligner technology introduced in 1997.⁴

Appliance Fabrication

AI-driven 3D scanning and CAD/CAM technology create custom-fitted orthodontic appliances (e.g., clear aligners). The development of digital clear aligner devices has made it easier to simulate orthodontic treatment, since manual methods made this procedure more difficult.²⁷ AI-generated data is fed into CAD (Computer-Aided Design) software to create digital models of braces, aligners, or retainers. CAD designs are then sent to CAM (Computer-Aided Manufacturing) systems, which use 3D printing or milling machines to create the appliance.²⁷ In 1946, Dr. Kesling introduced a removable plastic orthodontic appliance designed to make minor tooth movements—an innovation that laid the foundation for modern clear aligner therapy. Decades later, this concept evolved into a transformative orthodontic solution with the introduction of the Invisalign system by Align Technology in 1998.²⁶

Orthognathic Surgery Planning

AI models, especially those using facial images and key measurements, are increasingly used to support orthognathic surgery decisions by improving diagnostic accuracy and offering guidance, though challenges like feature selection and overfitting remain.⁹ AI is transforming orthognathic surgery by automating diagnostics, improving the accuracy of treatment planning, and enhancing surgical outcomes through tools like 3D simulations and machine learning-based decision systems.⁸

Cleft Lip and Palate (CLP)

The clinical application of AI in diagnosis and treatment planning of cleft patients can be seen from different aspects. **Diagnosis and Prediction:** Some studies used AI in identifying clefts and morphologic features in individuals with CLP. In this diagnosis ultrasound images can be used prenatally.¹⁴ Agarwal et al. used AI in the identification of digital camera images through Support Vector machine (SVM) with a previously trained convolutional neural network. This combination identified images of individuals with clefts from normal, and differentiated between unilateral and bilateral.¹⁴

Pre-surgical Orthopaedics: Sequential plates for infants with cleft lip and palate who required naso-alveolar moulding (NAM) was developed using AI. Moulding of foetal tissues was done, taking into account the growth of the maxillary arch using AI. A sequence of plates for NAM was generated, and this can be 3D-printed and delivered.¹⁴ In a study conducted among healthy control and non-syndromic cleft lip and palate infants (NSCL/P), machine learning algorithms were used in detecting defective variants in two genes necessary in folic acid and Vitamin A biosynthesis (methylenetetrahydrofolate reductase (MTHFR) and retinol-binding protein 4 (RBP4) genes).¹⁴ A study by Machado et al. on the Brazilian population for NSCL±P revealed interactions among 13 SNPs as well as defects in genes involved in folate metabolism.^{14,29}

Tele-orthodontics

Artificial intelligence (AI) has significantly advanced orthodontic remote monitoring by enhancing diagnosis, treatment planning, and patient follow-up. Systems like Dental Monitoring and Grin use AI-driven platforms to track oral hygiene, aligner progress, and treatment outcomes through patient-generated images and videos, reducing the need for in-office visits and improving efficiency³⁰. Complementing this, AI integration into telehealth systems, as demonstrated by Thurzo et al., has enabled real-time monitoring, early issue detection, and tailored interventions, improving patient compliance and allowing clinicians to manage more cases with personalised care⁴.

Benefits of AI in Orthodontics

- Faster and more accurate diagnoses
- Improved treatment planning
- Enhanced workflow and clinical efficiency
- Teleorthodontic capabilities
- Better patient experience and engagement
- Higher treatment success rates
- Significant time savings

Challenges and Limitations (^{4, 8, 28, 9})

- Data bias and quality issues
- High initial setup costs
- Lack of standardization and regulation
- Limited human oversight and ethical concerns
- Patient privacy and data security risks
- Resistance to change among practitioners
- Inaccuracy in complex or non-standard cases

Possibilities of Integration of AI in Underserved Countries

1. Fully Automated Diagnosis: Advanced AI models will be able to detect malocclusions, assess skeletal structures, and generate instant treatment plans with minimal human intervention. AI will integrate with CBCT, intraoral scanners, and panoramic X-rays to provide real-time orthodontic analysis. Automated detection of orthodontic treatment needs like the index of orthodontic

treatment need (IOTN) and index of orthognathic functional treatment need (IOFTN) would also be investigated.

2. 3D Printing Integration: AI will design and 3D-print fully customised clear aligners, braces, and orthodontic devices in real time. Machine learning algorithms will optimise instant tooth movement predictions, making treatments faster and more efficient. Patients may soon receive on-demand, in-office aligners without needing external manufacturers.

3. Smart Braces: AI-integrated smart braces will monitor tooth movement and automatically adjust forces using built-in sensors. Self-adjusting brackets with micro-robotics could gradually apply precise forces to reduce treatment time. AI-driven sensors will predict potential treatment delays and recommend adjustments in real-time.

4. AI + 5G for Remote Care: 5G networks will enable real-time AI analysis of patient scans from anywhere in the world. AI-powered remote monitoring tools will reduce clinic visits and allow orthodontists to track patient progress from their smartphones. Teleorthodontics will become more advanced, allowing orthodontists to provide instant consultations and treatment adjustments.

5. Virtual Assistants: AI chatbots will act as virtual assistants, answering patient queries, scheduling appointments, and providing treatment reminders. AI will analyse patient habits (e.g., aligner wear time)

and provide real-time compliance feedback. Patients will get personalised oral hygiene recommendations based on AI-driven risk assessments.

6. Predictive Analytics: AI will use big data analysis to predict treatment success rates and suggest the most efficient treatment plans. Machine learning models will forecast how teeth will move over time, allowing orthodontists to prevent complications before they arise. AI could also help in early orthodontic intervention for children, predicting future orthodontic issues before they fully develop.

7. Ethical Advancements: AI-driven orthodontics will require new regulations and ethical guidelines to ensure patient data privacy and accuracy. More orthodontists will undergo AI training, integrating AI seamlessly into daily practice.

Conclusion

Artificial Intelligence is reshaping orthodontics by enhancing diagnostics, personalising treatments, and streamlining workflows. With ongoing advancements, AI is set to make orthodontic care more precise, efficient, and accessible, ultimately leading to better outcomes and higher patient satisfaction

Authors' Contribution: All the authors contributed to the manuscript

Financial Support - Financed by the authors

Conflict of Interest - None to declare

References

1. Haenlein, M.; Kaplan, A. A Brief History of Artificial Intelligence: On the Past, Present, and Future of Artificial Intelligence. *Calif.Manag. Rev.* 2019, 61, 5–14.
2. Zhu X, Liu L, Sun Y, Li Y. Artificial Intelligence: A Powerful Paradigm for Scientific Research. *Trends in Science & Technology.* 2021;5(3):1-2. doi:10.1016/j.trst.2021.05.004.
3. Zaman BU. Big Data, Machine Learning, and AI in Decision Making. *ResearchGate.* Published October 2023. Available from: https://www.researchgate.net/publication/374616592_Big_Data_Machine_Learning_and_AI_in_Decision_Making
4. Amutha M, Kumaran NK. Artificial Intelligence in Orthodontics - A Narrative Review. *IOSR J Dent Med Sci.* 2024;23(8):38-43. doi:10.9790/0853-2308043843
5. Bichu YM, Hansa I, Bichu AY, Premjani P, Flores-Mir C, Vaid NR. Applications of artificial intelligence and machine learning in orthodontics: a scoping review. *Prog Orthod.* 2021;22(1):1-2.

6. Kazimierczak N, Kazimierczak W, Serafin Z, Nowicki P, Nożewski J, Janiszewska-Olszowska J. AI in Orthodontics: Revolutionizing Diagnostics and Treatment Planning—A Comprehensive Review. *J Clin Med*. 2024;13(2):1-19.
7. Carillo-Perez F, García-Godoy F, González-Cabezas C. Applications of artificial intelligence in dentistry: A comprehensive review. *J Esthet Restor Dent*. 2021;33(4):
8. Thakur SM, Shenoy U, Hazare A. et al. Transforming Orthodontics with artificial intelligence: A comprehensive review. *J Adv Dental Prac Res*. 2024;3(2):56-64.
9. Liu J, Chen Y, Li S, Zhao Z, Wu Z. Machine learning in orthodontics: Challenges and perspectives. *Adv Clin Exp Med*. 2021;30(10):1065–1074
10. Amasya H, Yildirim D, Aydogan T, Kemaloglu N, Orhan K. Cervical vertebral maturation assessment on lateral cephalometric radiographs using artificial intelligence: Comparison of machine learning classifier models. *Dentomaxillofac Radiol*. 2020;49(5):20190441.
11. Liu J, Zhang C, Shan Z. Application of artificial intelligence in orthodontics: current state and future perspectives. *Healthcare (Basel)*. 2023;11(20):2760. doi:10.3390/h
12. Dong W, Chen Y, Li A, Mei X, Yang Y. Automatic detection of adenoid hypertrophy on cone-beam computed tomography based on deep learning. *Am. J. Orthod. Dentofac. Orthop*. 2023;163, 553–560.e553.
13. Asiri Sn, Tadlock Lp, Schneiderman E, Buschang Ph. Applications Of Artificial Intelligence And Machine Learning In Orthodontics. *Apos Trends In Orthodontics*. 2020;10:17-24.
14. Dhillon H, Chaudhari PK, Dhingra K, Kuo R-F, Sokhi RK, Alam MK, Ahmad S. Current applications of artificial intelligence in cleft care: a scoping review. *Front Med*. 2021 Jul 28;8:676490. doi: 10.3389/fmed.2021.676490.
15. Hwang, H.W.; Park, J.H.; Moon, J.H.; Yu, Y.; Kim, H.; Her, S.B.; Srinivasan, G.; Aljanabi, M.N.A.; Donatelli, R.E.; Lee, S.J. Automated Identification of Cephalometric Landmarks: Part 2-Might It Be Better than Human? *Angle Orthod*. 2020, 90, 69–76.
16. Rudolph, D.J.; Sinclair, P.M.; Coggins, J.M. Automatic Computerized Radiographic Identification of Cephalometric Landmarks. *Am. J. Orthod. Dentofac. Orthop*. 1998, 113, 173–179.
17. Kök H, Acilar Am, İzgi Ms. Usage And Comparison Of Artificial Intelligence Algorithms For Determination Of Growth And Development By Cervical Vertebrae Stages In Orthodontics. *Prog Orthod*. 2019 Nov 15;20(1):41
18. Montúfar J, Romero M, Scougall-Vilchis Rj. Automatic 3-Dimensional Cephalometric Landmarking Based On Active Shape Models In Related Projections. *Am J Orthod Dentofacial Orthop*. 2018 Mar;153(3):449-458
19. Tomasetti P, Enz A, Roth C, Giunta RE, Ehrl D. F4CE: a mobile application for automated measurement of facial proportions using artificial intelligence. *Aesthetic Plast Surg*. 2024 Apr 4.
20. Jung Sk, Kim Tw. New Approach For The Diagnosis Of Extractions With Neural Network Machine Learning. *Am J Orthod Dentofacial Orthop*. 2016 Jan;149(1):127-33
21. Xie X, Wang L, Wang A. Artificial Neural Network Modeling For Deciding If Extractions Are Necessary Prior To Orthodontic Treatment. *Angle Orthod*. 2010 Mar;80(2):262-6.
22. Kumar V, Abdulkreem A, Tanmoy M, et al. Artificial intelligence-based automated preprocessing and classification of impacted maxillary canines in panoramic radiographs. *Dentomaxillofac Radiol*. 2024;53(1):20230289. doi:10.1259/dmfr.20230289.
23. Cadenas de Llano Perula M. AI-based prediction of maxillary canine eruption pathway [Internet]. KU Leuven Research Portal; 2024 Sep 2 [cited 2025 Apr 19]. Available from: <https://research.kuleuven.be/portal/en/project/3M240152>

24. Kivovics, M., Szanyi, S. M., Takács, A., Répási, M., Németh, O., & Mijiritsky, E. (2024). Computer-assisted open exposure of palatally impacted canines for orthodontic eruption: A randomized clinical trial. *Journal of Dentistry*, 147, 105110.
25. López-Castaño F, Arnabat-Domínguez J, Vela-Netto A, Gay-Escoda C. Artificial intelligence for orthodontic diagnosis and treatment planning. *Med Oral Patol Oral Cir Bucal*. 2024 Mar 1;29(2):e121–8..
26. Ruiz DC, Luz JG, Oliveira DD, Silva FFD, Elias CN. Unveiling the role of artificial intelligence applied to clear aligner therapy: A scoping review. *J Dent*. 2025 Mar;154:105564.
27. Smiles LA. Dental 3D printing in pediatric dentistry [Internet]. Los Angeles: Smiles LA; [cited 2025 Apr 21]. Available from: <https://smilesla.com/pediatric-dentistry-los-angeles/dental-3d-printing/>
28. Nordblom NF, Büttner M, Schwendicke F. Artificial Intelligence in Orthodontics: Critical Review. *J Dent Res*. 2024;103(6):577–584. doi:10.1177/00220345241235606.
29. Machado RA, de Oliveira Silva C, Martelli-Junior H, das Neves LT, Coletta RD. Machine learning in prediction of genetic risk of nonsyndromic oral clefts in the Brazilian population. *Clin Oral Invest*. (2021) 25:1273–80. doi: 10.1007/s00784-020-03433-y
30. Miranda F, Barone S, Gillot M, Baquero B, Anchling L, Hutin N, et al. Artificial intelligence applications in orthodontics. *J Calif Dent Assoc*. 2023;51(1):2195585. doi:10.1080/19424396.2023.2195585.

Talon Cusp: An uncommon occlusal anomaly – conservative management of two cases.

*Agboghroma GO^a, Akinyemi OA^a, Enabulele JE^b

Abstract

Talon cusp, a rare dental anomaly characterized by an extra cusp-like structure on anterior teeth, poses significant clinical challenges. This report presents two cases of talon cusp affecting maxillary central incisors, highlighting their clinical implications and successful conservative management.

The cases illustrate distinct complications, including premature contact, articulation difficulties, inadequate mouth closure, food impaction, and patient discomfort. A thorough literature review reveals variability in prevalence, etiology, and classification systems, underscoring the importance of radiographic evaluation and multidisciplinary approaches.

This report emphasizes the importance of early diagnosis and intervention to prevent complications and ensure optimal oral function and patient comfort. Conservative management strategies, including restorative and minor surgical interventions, are discussed, highlighting the potential for successful outcomes without extensive treatment.

Keywords: Talon Cusps, Problems, unilateral anomalous cups, conservative treatment

Authors' affiliate

^a Restorative Unit, Department of Dental and Maxillofacial Dentistry, National Hospital, Abuja.

^b Department of Restorative Dentistry, School of Dentistry, University of Benin.

Correspondence:

*Agboghroma GO

Email: okeoghenemaro@yahoo.com

Introduction

Talon Cusp, relatively rare developmental anomalies, present as accessory cusps resembling an eagle's talon, emanating from the cingulum area or cemento-enamel junction of anterior teeth, and so named for its semblance to an eagle's talon.^{1,2}

These anomalies can occur in both deciduous and permanent dentition. The etiology of talon cusps is unknown, but it is often attributed to the evagination of inner enamel epithelium during the morpho-differentiation stage of tooth development. It appears to be a balance between both environmental and genetic factors (Heaton and Pickering, 2013).

Prevalence varies considerably among ethnic groups.³ The variability in reported prevalence rates

of talon cusps may be attributed to the lack of standardised classification criteria and inconsistent definitions.⁴ To address this issue, Hattab et al. (1996) proposed a standardised talon cusp scoring system.⁵ This system categorises talon cusps into three distinct types based on their morphology:

Type 1: A prominent projection extending from the cingulum to more than half the distance to the incisal edge.

Type 2: A well-defined projection extending less than half the distance to the incisal edge.

Type 3: A minor expression characterised by enlarged cingula and their variations.

This classification system aims to provide a clear and consistent framework for diagnosing and reporting talon cusps, particularly those occurring on the lingual and palatal surfaces.

Talon cusps are composed of normal enamel, dentine, and may contain pulp tissue.⁶ They often attach to the incisal ridge, forming a T or Y shape, with no predilection for sex, although males are reportedly

more affected⁶. Once considered rare, radiographic examination has revealed a higher incidence.⁷ Their presentation is mostly unilateral and on the incisal towards the palatal with fewer bilateral or labial presentation⁵. Clinically, talon cusps are significant due to potential complications, including aesthetic concerns, speech difficulties, breastfeeding problems, attrition, occlusal interferences, temporomandibular joint pain, accidental cusp fracture, displacement of affected teeth, interference with tongue space, irritation of tongue during speech and mastication, periodontal problems due to excessive occlusal forces, development of dental caries in developmental groove of the talon cusps.⁴ This article presents two case reports of talon cusp affecting permanent teeth, highlighting diverse associated complications and successful conservative treatment outcomes.

Case 1

A 10-year-old male patient presented to the clinic accompanied by his mother, who expressed concerns regarding his dental aesthetics, tongue irritation, and difficulty closing his mouth, leading to mouth breathing and teeth grinding. The patient's deciduous teeth erupted and exfoliated normally. However, upon eruption of his permanent teeth, his mother noticed him frequently grinding his teeth and complaining of tongue irritation. As the issue persisted, she observed that he struggled to close his mouth adequately, prompting her to seek dental consultation.

The patient is the eldest of four siblings (one male and three females), with no family history of similar dental anomalies. His mother reported a normal pregnancy, without any complications or medication use during the pre- and postnatal periods. There was no history of trauma or associated systemic health issues.

On examination, the patient was a healthy and energetic young boy, presented with fully erupted

permanent dentition, excluding third molars. Soft tissue examination revealed minimal plaque accumulation on the palatal aspect of the maxillary central incisor and a slight diastema.

The patient exhibited a Class I molar relationship. Notably, both maxillary central incisors were slightly enlarged, with a prominent, horn-like projection palatally on the maxillary right central incisor (Figure 1). This talon cusp was large, well-defined, and hook-like, extending downward from the palatal surface of the crown, preventing adequate mouth closure.

The talon cusp exhibited a pointed, sharp tip with a slight curvature, causing tongue indentation. A moderate, non-carious dental groove was present at the junction of the talon cusp and the palatal surface of the tooth. Plaque accumulation and inflammation were observed around the marginal gingiva on the palatal surface of the maxillary right central incisor, eliciting slight bleeding on probing.

An occlusal radiograph confirmed the presence of a fully formed maxillary right central incisor, unaffected in its underlying structure (Figure 2). No additional dental anomalies or abnormalities were detected.

Treatment involved gentle reduction of the talon cusp using a high-speed turbine handpiece and diamond bur under copious water irrigation, without anesthesia, to assess sensitivity (Figure 3). Post-procedure, the patient was prescribed fluoride therapy and instructed to use fluoride toothpaste.

The patient was monitored closely through a series of follow-up appointments at weekly, bi-weekly, monthly, 3-month, and 6-month intervals, without any adverse reactions, such as color changes, pain, or periapical pathosis. Following an uneventful recovery, the patient was discharged.

A long-term follow-up examination four years later revealed that the tooth remained asymptomatic, indicating a successful treatment outcome.



Figure 1; Talon cusp on tooth 11



Figure 2; talon cusps on tooth 11



Figure 3; OPG showing talon cusp on tooth 11

Case 2

A 32-year-old female patient presented to the clinic with complaints of a projection and presence of an extra structure at the back of one of her teeth, along with intermittent pain originating from the affected tooth.

On examination, the patient was found to be a healthy young lady with a full complement of permanent dentition in a Class I molar relationship. The maxillary right lateral incisor appeared normal, with no discoloration. However, the palatal aspect revealed a horn-like projection attached to the cingulum, separated from the palatal surface by a small space that facilitated food entrapment and showed early signs of caries (Figure 4). Additionally, moderate cusp interference was observed, affecting normal occlusion.

The talon cusp was reduced through selective cuspal grinding utilising a flame-shaped diamond bur, accompanied by topical fluoride application. The patient was recalled monthly, and by the end of three visits, the cusp was virtually eliminated, although some sensitivity persisted (Figure 5). To ensure pulp protection, calcium hydroxide was applied, followed by a composite restoration.

The patient underwent an 8-month follow-up period, involving regular vitality testing and intra-oral radiographs. Throughout this time, the tooth remained asymptomatic with preserved vitality. Once stability was confirmed, the patient was discharged with instructions to return if any symptoms arose. To date, no further issues have been reported.



Figure 4: Talon cusp on tooth 12



Figure 5: periapical radiograph of Talon cusp on tooth 12

Discussion

Talon Cusp, also known as dens evaginatus, is a rare dental anomaly affecting anterior teeth. A systematic literature review and meta analysis reported a range of prevalence between 0.06 % and 40.8 %.¹⁴ Its etiology remains unknown, but it is attributed to disruptions during the morpho-differentiation stages of tooth development. Although associated with various syndromes, most reported cases occur in isolation, primarily in the permanent dentition.¹⁵ Notably, talon cusps are more prevalent in permanent teeth than deciduous ones.⁷

The talon cusp represents an extreme variation of normal tooth morphology, existing on a continuum that includes a normal cingulum, an enlarged cingulum (trace talon), a small accessory cusp (semi-talon), and a fully formed talon cusp.^{5,8}

The two reported cases of talon cusp presented unilaterally in the maxillary incisors, consistent with previous reports.^{15,16} However, the gender and age of the patients differed from previous studies. Our cases involved a male child and an adult female, whereas previous reports described female children¹⁵⁻¹⁷ and adults.^{5,6,17,18}

Notably, the location of the talon cusp varied between the two cases, affecting the right incisor in one and the left incisor in the other. This contrasts with a previous report where all cases occurred on the right side,¹⁵ but

aligns with another study that documented involvement of both right and left incisors.¹⁶

In agreement with some previous studies,¹⁵ both cases involved central incisors exclusively. However, this differs from reports describing talon cusps on lateral incisors only^{17,18} or both central and lateral incisors.¹⁶

Small talon cusps are typically asymptomatic and require no treatment. In contrast, larger talon cusps can pose significant challenges for both patients and clinicians, from diagnosis to treatment planning.⁹

The problems associated with talon cusps in the reported cases are consistent with previously documented issues.¹⁵⁻¹⁸ Patients with talon cusps may experience compromised aesthetics, breastfeeding difficulties, tongue irritation during speech or mastication, occlusal interference, and periapical and periodontal issues.^{2,5}

Dentists may face additional challenges, such as diagnostic difficulties, interference with orthodontic tooth movement, and increased risk of caries due to plaque and debris accumulation in the developmental groove.

The treatment of talon cusps requires meticulous planning, contingent upon the presence or absence of pulp tissue within the cusp. However, radiographic assessment is often hindered by the superimposition of the horn over the affected crown, and histologic examination may not reliably detect pulp tissue in the anomalous cusp.^{10,11}

In the reported cases, treatment was facilitated by the cusp's size and composition. The smaller cusp was successfully managed with grinding, without complications. The second tooth initially exhibited sensitivity, which resolved subsequently. This outcome aligns with previous reports where selective grinding and fluoride varnish application were effective.^{15,18}

Interestingly, the adult case presented similarly to paediatric cases, with comparable treatment outcomes. This contrasts with previous reports where adult patients required more extensive treatment, including grinding and root canal therapy.^{17,18}

Early diagnosis and timely treatment of talon cusps are crucial in preventing associated complications. Reports have highlighted the risks of anterior teeth subjected to shearing forces, leading to displacement of occluding teeth and potential fracture of the anomalous cusp.⁹

Prompt intervention is vital in maintaining the pulp-dentin complex's integrity, essential for preserving functional dentition. Selective grinding, fluoride varnish application, and sealant use can be enhanced by the natural formation of reparative dentine over time. Research indicates that reparative dentine formation occurs more rapidly in primary teeth than permanent teeth, and its thickness increases with

time, proportional to the remaining dentine thickness.

Studies^{12,13} provide valuable guidance on timed periods of selective grinding, emphasising the importance of close monitoring for color changes, vitality changes, and radiographic changes.

Early intervention and careful monitoring enable clinicians to mitigate potential complications and ensure optimal outcomes for patients with talon cusps.

Conclusion

Talon cusp is a rare developmental anomaly affecting anterior teeth, leading to varied clinical issues depending on its presentation. Prompt diagnosis and intervention are crucial in preventing complications. The two cases presented demonstrate the effectiveness of conservative treatment approaches, which were successfully employed to manage the anomaly. Therefore, conservative management is recommended for talon cusp cases, emphasising the importance of early detection and timely intervention.

Authors' Contribution: All the authors contributed to the manuscript

Financial Support - Financed by the authors

Conflict of Interest - None to declare

References

1. Hahab FN, Yasmin OM, Ai-Nimri KS. Talon Cusp- Clinical significance and management: Case reports. *Quint Int*;26: 115-20,1995
2. Mellor JK, Ripa LW. Talon cusp: a clinically significant anomaly. *Oral Surg Oral Med Oral Pathol*. 1970 Feb;29(2):225-8. doi: 10.1016/0030-4220(70)90089-7. PMID: 5262843.
3. Sedano HO, Freyne IC, Garza de la Garza NL et al. (1989) Clinical Orofacial Abnormalities in Mexican Children. *Oral Surgery, Oral Medicine, Oral Pathology* 68, 300-11
4. Al-Omari MA, Hattab FN, Darwazeh AM, Dummer PM. Clinical problems associated with unusual cases of talon cusp. *Int Endod J*. 1999 May;32(3):183-90. doi: 10.1046/j.1365-2591.1999.00212.x. PMID: 10530205.
5. Hattab FN, Yassin OM, al-Nimri KS. Talon cusp in permanent dentition associated with other dental anomalies: review of literature and reports of seven cases. *ASDC J Dent Child*. 1996 Sep-Oct;63(5):368-76. PMID: 8958353.
6. Hattab FN, Yassin OM. Bilateral talon cusps on primary central incisors: a case report. *Int J Paediatr Dent*. 1996 Sep;6(3):191-5. doi: 10.1111/j.1365-263x.1996.tb00241.x. PMID: 9115977.
7. Tulunoglu O, Cankala DU, Ozdemir RC. Talon's cusp: report of four unusual cases. *J Indian Soc Pedod Prev Dent*. 2007 Mar;25(1):52-5. doi: 10.4103/0970-4388.31993. PMID: 17456971.
8. Davis, P., Brook, A. The presentation of talon

- cuspid: diagnosis, clinical features, associations and possible aetiology. *Br Dent J* **160**, 84–88 (1986).
9. M.A.OAI-Omari, F.N Hattab² AMG Darwazeh³ & P.M.H Dummer.⁴ Clinical Problems Associated with Unusual Cases Of Talon Cusp. *International Endodontic Journal* 1999 **32**, 183-190.
 10. Mader & Kellog 1985. Primary Talon Cusp. *Journal of Dentistry For Children* **52**,223-6.
 11. Salama FS, Hanes CM, Hanes PJ. Ready MA (1990) Talon Cusp: A Review and Two Case Reports on Supernumerary Primary and Permanent Teeth. *Journal of Dentistry For Children* **57**,147-50.
 12. Scotti R. Villa L. Carossa S. Clinical Applicability of The Radiographic Method For Determining The Thickness of Calcified Crown Tissues. *J.Prosthet Dent.* 1991 **65**: 65-7
 13. Seltzer, Bendois. *The Dental Pulp, Biologic Considerations in Dental Procedure.* 3 ed. Chennai. All India Publishers and Distributors 2002 P.324-48.
 14. Pierre-Hadrien Decaup, Elsa Garot, Patrick Rouas, Prevalence of talon cusp: Systematic literature review, meta-analysis and new scoring system, *Archives of Oral Biology*, 2021;**125**: 105112,ISSN 0003-9969,
 15. Bansal AV, Choudhary P, Kulkarni VK, Bansal A, Shashikiran ND. Talon cusps: conservative management. *J Clin Pediatr Dent* 2011;**35**:345-348.
 16. Butt R,Aspinall A, Tanday A, Brown CJ. Talon cusp management: A case series. *Dental Update* 2024;**49**:707-709.
 17. Ozcelik B, Atila B. Bilateral palatal talon cusps on permanent maxillary lateral incisors: a case report. *Eur J Dent.* 2011 Jan;**5**(1):113-6. PMID: 21228961; PMCID: PMC3019756.
 18. Ozer SY. Talon Cusp: A literature review and three case reports. *J Int Dent Med Res* 2009;**2**:45-49.



Nigerian Association of Orthodontists

19th

**Annual General Meeting/
Scientific Conference**



Main theme:

**Embracing technology and innovations
in Orthodontics: Pioneering the future**

Sub theme:

**The business of Orthodontics and its long term
commitment: Planning retirement and succession.**

 **23rd - 26th September 2025**
 **8am Daily**

 **View Points Hotel and JOOPSA
Benin City Edo State.**

Instructions for Authors

West African Journal of Orthodontics is a peer-reviewed journal published by affiliated Orthodontic Groups and Associations in the West African Sub region. The journal gives priority to reports of outstanding clinical and experimental and epidemiological works on malocclusion, dento-facial defects as well as important contributions related to common orthodontic problems in children, adolescents and adults worldwide.

Submission

Manuscripts and registered letters should be sent to: the Editor, West African Journal of Orthodontics, Department of Child Dental Health, Faculty of Dentistry, College of Health Sciences Obafemi Awolowo University, Ile-Ife, Osun State. Nigeria.

Manuscripts in MS word attachments may also be submitted via Email to wajoeditorinchief@yahoo.com, in addition to hard copies. Tables, figures and text should be included in the same file if possible. Authors may submit their research works by email only; such manuscripts need not be simultaneously sent by post.

However, photographs and/or figures need to be sent separately as hard copy (under figures and illustrations).

Acceptance

Manuscripts should meet the following criteria: original material, clear writing, appropriate study methods, valid data, and reasonable conclusions supported by the data, in short, they should contain important information on topic of general orthodontic interest.

Peer-review Process

All the manuscripts that adhere to its style and Instructions for Authors are referred to peer-review. Some of them are rejected immediately after an inhouse review. The rejection at this stage is due to insufficient originality, serious scientific flaws or absence of message. The remaining articles are sent to at least two reviewers who are experts in the subject. Manuscripts are reviewed with due respect for authors' confidentiality, and the identity of peer reviewers is also kept confidential. A decision is made from 6 to 12

weeks according to the response from reviewers, revision by the author(s) and reappraisal on the revision.

The accepted manuscripts are subjected to editorial revision to comply with the requirements on language and style of the journal. The rejected manuscript is not returned to authors but its copies are kept for 3 months to answer any queries. The copyright of the accepted and published articles is held by the journal and all the published materials cannot be reproduced or published elsewhere, in whole or part, without the written permission from the editor.

Duplicate Submission

Manuscripts are considered with the understanding that they have not been published previously and are not under consideration by another publication. The author should alert the editor if the work includes subjects about which a previous report has been published. A research paper submitted to this journal should not overlap by more than 10% with the previously published material or work submitted elsewhere, which would be considered as duplicate publication. If in doubt, authors may forward copies of the published work or material submitted elsewhere to this journal for decision making.

Proofs and Reprints

The corresponding author of the accepted article shall be supplied with the proof. Corrections on the proof should be restricted to errors only and no substantial additions/deletions should be made. No addition or deletion in the names of the authors is permissible at this stage. A copy of the issue carrying the article is supplied free of charge to the authors.

Reprints may be ordered on payment in advance.

Categories of Articles

Articles can be sent as editorials, original articles, review articles, special communications, brief reports, case reports, letters to editor, commentaries, or for images section.

They are mostly included under Events of Interest free of cost. This journal reserves the right to be selective in publishing these announcements.

Preparing Manuscripts

Manuscripts should be prepared in accordance with the Uniform Requirements for Manuscripts submitted to Biomedical Journals. 2 A summary of technical requirements for preparing the manuscript is provided below:

- Three copies of the manuscript should be submitted.
- Use 1 side of standard size 21.6x27.9 cm A4, white bond paper, with margins of at least 2.5 cm on each side.
- Double-space throughout including title page, abstract, text, acknowledgements, references, tables and figure legends. Start each of these sections (in same order) on a new page, numbered consecutively in the upper right hand corner, beginning with the title page.
- Use at least 12 point font size (Times New Roman or Arial).
- Submit photographs and transparencies in a separate heavy paper envelope (enclosed in cardboard, to prevent bending during mail handling).
- Conventional units are preferred with SI units in parenthesis, if available. The metric system is preferred for the expression of length, area, mass and volume.
- Use nonproprietary names of material rugs, devices and other products.
- All manuscripts should be accompanied by a signed statement by all authors regarding authorship, responsibility, financial disclosure and acknowledgements, as per standard format (Appendix J)[23 1 Those sending their manuscript through email are also required to submit this form by post with original signatures.

Manuscripts not fulfilling the technical requirements shall be returned to the authors without initiating the peer-review process.

Title Page

The page should contain (i) the title of the article: which should be concise but informative (simpler the title the better; preferably it should contain all the key words to help electronic retrieval reliably); (ii) a short

running title of less than 40 characters placed at the foot end of the title page; (iii) initials and surname of each author with the highest academic degree(s) and designation at the time when the work was done; (iv) details of the contribution of each author; (v) name of department(s) and institution(s) to which the work should be attributed; (vi) disclaimers, if any; (vii) name, address, telephone, fax, email address of the corresponding author, (viii) source(s) of support in the form of grants, equipment, drugs or all of these; and (ix) declaration on competing interests.

Authorship

All persons designated as authors should qualify for the authorship. Authorship credit should be based on substantial contributions to (i) concept and design, or acquisition of data, or analysis and interpretation of data; (ii) drafting the article or revising it critically for important intellectual content; and (iii) final approval of the version to be published. Conditions 1, 2 and 3 must all be met. Participation solely in the acquisition of funding or the collection of data does not justify authorship. All such people who contributed to the work but do not satisfy all the conditions should be listed in the acknowledgements.

Authors are responsible for obtaining written permissions from everyone acknowledged by name. One of the authors shall act as guarantor of the paper and he/she should take the responsibility for the integrity of the work as a whole, from its inception to published article.

Authors should provide a description of what each author contributed on the title page. Subsequently, no names can be added or deleted without written permission of the editor. Written consent of authors whose names are being deleted should be obtained.

This journal reserves the right to satisfy itself regarding the specific role of each listed author to justify authorship. All authors must give signed consent to publication (Appendix 1).

Competing Interest

Competing interest for a given manuscript exists when the author has ties to activities that could inappropriately influence his or her judgment, whether or not judgment is in fact affected. Financial relationships with industry for example, through employment, consultancies, stock ownership, honoraria, expert testimony, either directly or through immediate family, are usually considered to be the most important competing interests. However, conflicts can

Original Article

Original articles should report original research relevant to basic and clinical orthodontics including randomized trials, intervention studies, studies of screening and diagnostic tests, cohort studies, cost effectiveness analyses and case control studies. While reporting randomized controlled trials (RCT), authors must attempt to be in conformity with the consolidated standards of reporting trial.

(CONSORT) statements

Each manuscript should be accompanied with a structured abstract (divided into background, methods, results and conclusions) in no more than 250 words. Four to five key words to facilitate indexing should be provided in alphabetical order along with the abstract. The text should be divided in sections on introduction, methods, results, discussion and conclusion.

Acknowledgment section may be included where necessary. Number of tables and figures should be limited to the very relevant ones and may be compressed if necessary. The typical text length for such contributions is 2500-3 500 words (excluding title page, abstract, tables, figures, acknowledgments and references).

Brief Report

Short accounts of original studies are published as brief reports. The text should be divided into sections, i.e., abstract, introduction, methods, results and discussion.

Abstract should be of 100-150 words highlighting the aims, methods and main results along with 3-4 key words.

The text should contain no more than 1500 words, 3 illustrations or tables and up to 20 references, preferably recent publications.

Review Article

State-of-the-art review articles or systematic, critical assessments of literature are also published. Normally a review article on a subject already published in the West African Journal of Orthodontics is not accepted for a period of 3 years.

The typical length for review articles is 2000-3000 words, excluding tables, figures, and references.

Authors submitting review manuscripts should include a structured abstract of around 200 words describing the need and purpose of review, methods used for selection, extraction and synthesis of data, and main conclusions.

Clinical cases highlighting uncommon malocclusion condition, orthodontic treatment techniques are published as case reports. Single case reports are usually not accepted, unless some new or unusual aspect regarding aetiopathogenesis, diagnosis or management is brought out that adds to the existing body of knowledge. The text should not exceed 1000 words and is divided into sections, i.e., abstract, introduction, case report and discussion. The number of tables/figures should be limited to 2. Ten recent references are acceptable. A maximum of 3 or 1 author is permitted from the principle and each of the associated departments respectively. Thus, case reports from only one investigative department can have a maximum of 3 authors.

Letter to Editor(s)

Letters commenting upon a recent article in the West African Journal of Orthodontics are welcome.

Such letters should be received within 6 months of the article's publication. At the editorial board's discretion, a letter may be sent to authors! experts for comments and both letter and reply may be published together. Letters may also relate to other topics of interest to orthodontists and others, and/or useful clinical observations. Letters should not be more than 400 words. The number of authors should not exceed 2, including the authors' reply in response to a letter commenting upon an article published in this journal.

Images Section

A short text of about 150 words depicting the condition with color photographs (vide infra) is needed.

Normally only clinical photographs are accepted but accompanying skiagrams or pathological images could also be considered for publication.

Photographs should be of high quality, clearly identify the condition and preferably add to the existing knowledge.

Personal Viewpoint

Such articles are published on topical orthodontic issues including social aspects. It is expected that the authors have sufficient credible experience on the subject for giving viewpoints. These should not exceed 1500 words.

Notes, News and Events of Interest

Announcements for conferences, symposia, meetings or courses may be sent for publication in advance. The announcements should provide title, date(s) and place of the event and contact address, telephone, and email

occur for other reasons, such as personal relationships, academic competition and intellectual passion. If any of the authors have accepted reimbursement for attending symposium, a fee for speaking, fee for organizing educational reach, funds for a member of the staff of consultation fees from an organization that may in: way gain or lose financially from the result of the study, review, editorial or letter, a competing interest would be deemed to exist. If any of the authors had been employed by an organization that may in any way gain or lose financially from the publication, or if any of them hold stocks or shares in such an organization, competing interest would be deemed to exist. If competing interest exists, the author(s) must disclose them while submitting the manuscript.

Abstract and Key Words

The second page should carry an abstract in case of original article (250 words), review article (200 words), brief report (100-150 words), and case report (50 words), respectively. For original article and reviews, the abstract should be structured as detailed earlier. For brief reports, the abstract should state the purpose of the study, basic methodology, main findings (giving specific data and statistical significance) and key conclusion(s). Below the abstract, authors should provide 3-5 key words for indexing; terms from the Medical Subject Headings (MESH) list of Index Medicus should be used. The basic structure of a paper follows the well known acronym IMRAD, which stands for Introduction (what questions was asked), Methods (how was it studied), Results (what was found) and Discussion⁴.

Introduction

The introduction must clearly state the question that the author(s) tried to answer in the study. It may be necessary to briefly review the relevant literature. Only cite those references that are essential to justify the proposed study.

Materials and Methods

The methods section should describe, in a logical sequence, how the study was designed (e.g., how randomization was done), carried out (e.g., how subjects were chosen or excluded, ethical considerations, accurate details of materials used, exact drug dosage and form of treatment, etc.) and data were analyzed (e.g., an estimate of the power of the study, exact test used for statistical analysis, etc.). For standard methods, appropriate references are sufficient, but if standard methods are modified these should be clearly brought out.

Authors should provide complete details of any new methods or apparatus used (manufacturer's name and address in parentheses).

Ethics

When reporting experiments on human subjects, authors should indicate whether the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional or regional) and with the Helsinki Declaration of 1964, as revised in 2000.

They should indicate whether the study was approved by the Institutions' Ethical Committee, and whether informed consent was obtained from the study participants. They should not use patients' names, initials, or hospital numbers, especially in illustrative material. This journal reserves the right to reject a manuscript on ethical grounds, on the basis of recommendations of its "Ethical Committee", even if the research has been cleared by the institutional ethical committee. Moreover, when reporting experiments on animals, authors should indicate whether the institutional and national guide for the care and use of laboratory animals was followed.

Statistics

Authors should describe statistical methods with enough detail to enable a knowledgeable reader with access to the original data to verify the reported results. When possible, they meet to quantify findings and present them with appropriate indicators of measurement error or uncertainty (such as confidence intervals). Actual P values are provided rather than stating as just <0.05 or >0.05 etc. References for the design of the study and statistical methods should be to standard works when possible (with pages stated) rather than to papers in which the designs or methods were originally reported. Any general-use computer programs used should be specified and statistical terms, abbreviations, and most symbols be defined.

Results

This section should include only relevant, representative data and not all information collected during the study. Major findings should be presented clearly and concisely. Text, tables, and illustrations should be used sensibly while avoiding repeating in the text all the data depicted in the tables or illustrations and emphasizing or summarizing only important observations. Tables and figures should be restricted to those needed to explain the argument of the paper and to assess its support. It is necessary to cite the tables in the text and type them on separate sheets. It may also be useful to mention what the study did not find.

Discussion

Discussion ordinarily should not be more than one third of the total length of the manuscript. This section should include a summary of the major findings, their relationship to other similar studies, limitations of methods and implications of these findings in future research. Conclusions should be linked to the goals of the study. Unqualified statements and conclusions which are not completely supported by the data should be avoided. Authors should also refrain from making statements on economic benefits and costs unless their manuscript includes economic data and analyses.

Acknowledgements

In acknowledgements section, it is suitable to list all contributors who do not meet the criteria for authorship, such as a person who provided purely technical help, writing assistance, or a department head who provided only general support. Financial and material support should also be acknowledged.

Groups of persons who have contributed materially to the paper but whose contributions do not justify authorship may be listed under a heading such as "clinical investigators" or "participating investigators", and their function or contribution should be described, for example, "served as scientific advisers", "critically reviewed the study proposal", "collected data", or "provided and cared for study patients". A written consent is required from all the persons acknowledged, indicating their acceptance for the same.

Contributions to joint-authorship

In the case of multiple author-ship, authors are expected to state clearly their contributions to the paper being considered for publication in terms of study initiation, design including methodology, data collection, analysis and final write-up. The editorial board reserves the right to remove any author's name if the contribution is insignificant.

References

References should be numbered consecutively in the order in which they are first mentioned in the text.

References are identified in text, tables, and legends by Arabic numerals in parentheses. References cited only in tables or in legends to figures should be numbered in accordance with the sequence established by the first identification in the text of the particular table or figure.

The titles of journals should be abbreviated according to the style used in Index Medicus. Authors are required not to use abstracts, unpublished observations and personal communications as references. References to papers accepted but not yet published should be designated as "in press"; authors should obtain written permission to cite such papers as well as verification that they have been accepted for publication.

The references must be verified by the author against the original documents. The Uniform Requirements style (the Vancouver style) is based largely on an American National Standards Institute (ANSI) standard style adapted by the NLM for its databases.

Journal Article

List all authors when 6 or less. When 7 or more, list only first six and add et al. Ngan P, Yiu C, Hu A, Hagg U, Ei SHY, Gunel E. Cephalometric and occlusal changes following maxillary expansion and protraction. *Eur J Orthod* 1998; 20: 237-254.

Organization as Author

Australian Dental Association Inc. An Australian Schedule of Dental Services and Glossary. 7th edn. Sydney: Australian Dental Association Inc., 1996.

Complete Book

Department of Health. Shifting the balance of power within the NHS: securing delivery. London: Doll, 2001.

Clayton D, Hills M. Statistical models in epidemiology. Oxford: Oxford University Press, 1993.

Farkas LG. Anthropometry of the Head and Face, 2nd Edn, New York; Raven Press; 1994

Book Chapter Lekholm U, Zarb GA. Patient selection and preparation. In: Branemark P1, Zarb GA, Albrektsson T, editors.

Tissue integrated Prostheses: Osseointegration in Clinical Dentistry, Chicago: Quintessence; 1988,199-209

Thesis and Dissertation

Yong SJ. Bone mineral density of normal Korean adults. Ph.D. Thesis. Seoul, Korea; 1989 Anozike, AN. Orthodontic treatment needs and its impact on oral health related quality of life in Lagos school children aged 12-16 years. FMCDs. Dissertation. Lagos, Nigeria; 2006

Conference Proceedings

Marshall SJ, Rixon RC, Whiteford DN, Cumming JT. The OrthoForm 3-Dimensional Clinical Facial Imaging System. Proceedings of the 15th IFHE Congress 1998; 15:83-87.

Dictionary and Similar References

Stedman's medical dictionary. 26th ed. Baltimore: Williams & Wilkins; 1995. Apraxia; p.11 9-120. Unpublished accepted material Leshner Al. Molecular mechanism of cocaine addiction. N Eng J Med. In Press 1996.

Material from Internet

World Health Organization, 2002.
www.who.int/mental-health/prevention/suicide
(accessed August 1, 2004).

Tables

Each table should be typed in double-space on a separate sheet of paper. Tables not submitted as photographs must be numbered consecutively (Arabic numerals) in the order of their first citation in the text, with a brief but self explanatory title for each.

Each column should have a short or abbreviated heading. Explanatory matters are placed in footnotes, not in the heading. In footnotes all nonstandard abbreviations that are used in each table should be explained adequately. Statistical measures of variations should be identified such as standard deviation and standard error of the mean. Be sure that each table is cited in the text. If data are used from another published or unpublished source, it is necessary to obtain permission and acknowledge them fully.

Figures and Instructions

Figures should be professionally drawn and photographed; freehand or typewritten lettering is unacceptable. Instead of original drawings, X-ray films, and other material, sharp, glossy, black-and-white photographic prints of high quality are necessary, usually 127x 173 mm (5x7 in) but no larger than 203x254 mm (8x10 in) For color illustrations negatives or positive transparencies are provided, along with color prints. It is preferable to have the photograph in portrait form rather than in landscape form to fit easily into one column. Letters, numbers and symbols in photographs should be clearly legible.

Each figure should have a label pasted on its back indicating the number of the figure, author's name, and an arrow to mark the top and left side of the figure.

It is unacceptable to write on the back of figures or scratch or mark them by using paper clips, and to bend figures or mount them on cardboard. If photographs of individual/people are used, either the subjects must not be identifiable or their pictures must be accompanied by written permission to use the photograph. It is advisable to cover the eyes unless specifically need to be shown. If a figure has been published, the original source should be acknowledged and written permission from the copyright holder be obtained to reproduce the material. Figures should be numbered consecutively (Arabic numerals) according to the order in which they have been first cited in the text.

Legends for Illustrations

Legends for illustrations should be typed or printed out in double-space, starting on a separate page, with Arabic numerals corresponding to the illustrations.

When symbols, arrows, numbers, or letters are used to identify parts of the illustrations, each of them must be identified and explained in the legend. The internal scale should be explained and the method of staining in photomicrographs be identified.

Units of Measurement

Measurements of length, height, weight, and volume should be reported in metric units, i.e., meter(m), gram(g), or liter(l) or their decimal multiples.

Milliliter or deciliter should be expressed as ml or dl.

Red and white blood cell counts are to be expressed as $63 \times 10^6 / \text{mc l}$ and $\times 10^6 / \text{mc}$ respectively. Temperatures should be given in degrees Celsius and blood pressures in millimeters of mercury (mmHg). All hematological and clinical chemistry measurements should be reported in the conventional system or in terms of the International System of Units (SI).

Abbreviations and symbols

Only standard abbreviations are used in the text while avoiding abbreviations in the title and abstract.

The full term for which an abbreviation stands should precede its first use in the text unless it is a standard unit of measurement. Year, month, day, hour, minute and second should be abbreviated as yr, mon, d, h, mm, and s in tables respectively.

References

1. Mother M, Schulz KF, Altman DG, for the CONSORT Group. The CONSORT statement Revised recommendations for improving the quality of reports of parallel group randomize Trials. Lancet 2001; 357: 1191-1194. (Also available from: URL: <http://www.consort-statement.org/>). Accessed June 28, 2002.
2. International Committee of Medical Journal Editors. Uniform Requirements for Manuscripts Submitted to Biomedical Journals. Ann Intern Med 1997;126:36-47. (Updated October 2001 version Available from: URL: <http://www.icmje.org/>). Accessed June 28,2002.
3. JAMA Instructions for Authors. Available from URL: <http://jama.ama-assn.org/>. Accessed June 28, 2002.
4. Hall GM. Structure of a scientific paper. In: Hall GM, ds. How to write a paper. London:BMJ Books, 2000.
5. 52nd WMA General Assembly. World Medical Association Declaration of Helsinki. Ethical principles for medical research involving human subjects. Available from: URL: <http://www.wma.net/>. Accessed June 28,2002.

Appendix 1:

Declaration of Originality and Transfer of Copyright

(Please download from Nigerian Association of Orthodontists (NAO) website <https://www.nao-ng.org/>)

This form is to be submitted with the initial copies of the manuscript to: West African Journal of Orthodontics, Department of Child Dental Health, Obafemi Awolowo University Ile-Ife, Osun State. Nigeria Manuscript No. (If known):

The author(s) hereby affirms that the submitted manuscript entitled:

I/We certify that the manuscript represents valid work and that neither this manuscript nor one with substantially similar content under my/our authorship has been published or is being considered for publication elsewhere. For papers with more than I author, we agree to allow the corresponding author to serve as the primary correspondent with the editorial office, to review the edited typescript and proof.

I/We have seen and approved the submitted manuscript. All of us have participated sufficiently in the work to take public responsibility for the contents. All the authors have made substantial contributions to the intellectual content of the paper and fulfill at least 1 condition for each of the 3 categories of contributions: i.e., Category 1 (conception and design, acquisition of data, analysis and interpretation of data), Category 2 (drafting of the manuscript, critical revision of the manuscript for important intellectual content) and Category 3 (final approval of the version to be published).

I/We also certify that all my/our affiliations with or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript are completely disclosed on the title page of the manuscript. My/our right to examine, analyze, and publish the data is not infringed upon by any contractual agreement.

I/We certify that all persons who have made substantial contributions to the work reported in this manuscript (e.g., data collection, writing or editing assistance) but who do not fulfill the authorship criteria are named along with their specific contributions in an acknowledgment section in the manuscript. If an acknowledgment section is not included, no other persons have made substantial contributions to this manuscript.

I/We also certify that all persons named in the acknowledgment section have provided written permission to be named.

The author(s) undersigned hereby transfer(s), assign(s), or otherwise convey(s) all copyright ownership, including any and all rights incidental thereto, exclusively to the West African Journal of Orthodontics, in the event that such work is published in the West African Journal of Orthodontics.

Authors name(s) in order of appearance in the manuscript; signatures (date):

